



**QUEEN'S
UNIVERSITY
BELFAST**

Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey: participant-level analysis of two large prevention studies

Barbui, C., Tedeschi, F., Acarturk, C., Anttila, M., Au, T., Baumgartner, J., Carswell, K., Churchill, R., Cuijpers, P., Karyotaki, E., Klein, T., Koesters, M., Lantta, T., Nosè, M., Ostuzzi, G., Pasquini, M., Prina, E., Sijbrandij, M., Tarsitani, L., ... Purgato, M. (2022). Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey: participant-level analysis of two large prevention studies. *International Journal of Social Psychiatry*.

Published in:

International Journal of Social Psychiatry

Document Version:

Peer reviewed version

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2022 The Authors.

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey: participant-level analysis of two large prevention studies

Journal:	<i>International Journal of Social Psychiatry</i>
Manuscript ID	ISP-22-0527.R1
Manuscript Type:	Original Research Article
Date Submitted by the Author:	22-Sep-2022
Complete List of Authors:	<p>Barbui, Corrado; University of Verona, Tedeschi, Federico; University of Verona Acarturk, Ceren ; Koc University College of Social Sciences and Humanities, Psychology Anttila, Minna; University of Turku Au, Teresa; World Health Organization Baumgartner, Josef; University of Vienna Carswell, Kenneth; World Health Organization Churchill, Rachel; University of York Cuijpers, Pim; VU-University, Faculty of Clinical Psychology Karyotaki, Eirini; Vrije Universiteit Amsterdam Klein, Thomas; University of Ulm and BKH Günzburg, Department of Psychiatry II Kösters, Markus; Universitätsklinikum Ulm Klinik für Psychiatrie und Psychotherapie II Günzburg Lantta, Tella; University of Turku Nosè, Michela; University of Verona Ostuzzi, Giovanni; University of Verona Pasquini, Massimo; Policlinico Umberto I, Sapienza University of Rome, Department of Human Neurosciences Prina, Eleonora; University of Verona Sijbrandij, Marit; Vrije Universiteit Amsterdam Tarsitani, Lorenzo; University of Rome La Sapienza, Turrini, Giulia; University of Verona Uygun, Ersin; Istanbul Bilgi University Valimaki , Maritta ; University of Turku, Department of Nursing Science, University of Turku Walker, Lauren; University of York Wancata, Johannes; University of Vienna White, Ross; Queen's University Purgato, Marianna; University of Verona</p>
Keywords:	migrants, asylum seekers, refugees
Abstract:	<p>Aim: This study investigated the risk factors for mental disorder development in a large group of asylum seekers and refugees resettled in high- and middle-income settings.</p> <p>Methods: Participant-level data from two randomized prevention studies involving asylum seekers and refugees resettled in Western European countries and in Turkey were pooled. The two studies randomized participants with psychological distress, but without a diagnosis of mental disorder, to the Self-Help Plus psychological intervention or</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>enhanced care as usual. At baseline, exposure to potentially traumatic events was measured using the Harvard Trauma Questionnaire-part I, while psychological distress and depressive symptoms were assessed with the General Health Questionnaire and the Patient Health Questionnaire. After three and six months of follow-up, the proportion of participants who developed a mental disorder was calculated using the Mini International Neuropsychiatric Interview.</p> <p>Results: A total of 1,101 participants were included in the analysis. At three- and six-month follow-up the observed frequency of mental disorders was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. After controlling for confounders, logistic regression analysis showed that participants with a lower education level ($p = 0.034$), a shorter duration of journey ($p = 0.057$), and arriving from countries with war-related contexts ($p = 0.017$), were more at risk of developing mental disorders. Psychological distress ($p = 0.004$), depression ($p = 0.001$), and exposure to potentially traumatic events ($p = 0.020$) were predictors of mental disorder development.</p> <p>Conclusions: This study identified several risk factors for the development of mental disorders in asylum seekers and refugees, some of which may be the target of risk reduction policies. The identification of asylum seekers and refugees at increased risk of mental disorders should guide the implementation of focused preventative psychological interventions</p>



1
2
3
4 **Risk factors for mental disorder development in asylum seekers and**
5 **refugees resettled in Western Europe and Turkey: participant-level**
6 **analysis of two large prevention studies**
7
8
9
10
11
12

13 Corrado Barbui¹, Federico Tedeschi¹, Ceren Acarturk², Minna Anttila³, Teresa Au⁴, Josef
14 Baumgartner⁵, Ken Carswell⁴, Rachel Churchill⁶, Pim Cuijpers^{7,8}, Eirini Karyotaki⁷, Thomas
15 Klein⁹, Markus Koesters⁹, Tella Lantta³, Michela Nosè¹, Giovanni Ostuzzi¹, Massimo Pasquini¹⁰,
16 Eleonora Prina¹, Marit Sijbrandij⁷, Lorenzo Tarsitani¹⁰, Giulia Turrini¹, Ersin Uygun¹¹, Maritta
17 Välimäki^{3,12}, Lauren Walker¹³, Johannes Wancata⁵, Ross G. White¹⁴, Marianna Purgato¹
18
19
20
21
22

23 ¹ WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation,
24 Department of Neuroscience, Biomedicine and Movement Sciences, Section of Psychiatry,
25 University of Verona, Verona 37134, Italy
26

27 ² Department of Psychology, Koc University, Istanbul, Turkey
28

29 ³ University of Turku, Department of Nursing Science, 20014 University of Turku, Finland
30

31 ⁴ Department of Mental Health and Substance Use, World Health Organization, Geneva,
32 Switzerland
33

34 ⁵ Clinical Division of Social Psychiatry, Department of Psychiatry and Psychotherapy, Medical
35 University of Vienna, Vienna, Austria
36

37 ⁶ Centre for Reviews and Dissemination, University of York, York, UK
38

39 ⁷ Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health
40 research institute, Vrije Universiteit Amsterdam, WHO Collaborating Centre for Research and
41 Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam
42

43 ⁸ Babeş-Bolyai University, International Institute for Psychotherapy, Cluj-Napoca, Romania
44

45 ⁹ Department of Psychiatry II, Ulm University, Ulm, Germany
46

47 ¹⁰ Department of Human Neurosciences, Sapienza University of Rome, Italy
48

49 ¹¹ Trauma and Disaster Mental Health, Istanbul Bilgi University, Eyüpsultan/Istanbul, Turkey
50

51 ¹² Xiangya Nursing School, Central South University, Xiangya Center for Evidence-Based Practice
52 and Healthcare Innovation, 172 Tongzipo Road, Changsha, Hunan 410013, China
53

54 ¹³ Department of Health Sciences, University of York, York, UK
55

56 ¹⁴ School of Psychology, Queen's University Belfast, Belfast, UK
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Correspondence to:

Professor Corrado Barbui, WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, Ospedale Policlinico GB Rossi, Piazzale Scuro 10, 37100 Verona, Italy. Email: corrado.barbui@univr.it

Word count: 4,472

For Peer Review

Abstract

Background: In asylum seekers and refugees, the frequency of mental disorders, such as depression, anxiety, and post-traumatic stress disorder, is higher than the general population, but there is a lack of data on risk factors for the development of mental disorders in this population.

Aim: This study investigated the risk factors for mental disorder development in a large group of asylum seekers and refugees resettled in high- and middle-income settings.

Methods: Participant-level data from two randomized prevention studies involving asylum seekers and refugees resettled in Western European countries and in Turkey were pooled. The two studies randomized participants with psychological distress, but without a diagnosis of mental disorder, to the Self-Help Plus psychological intervention or enhanced care as usual. At baseline, exposure to potentially traumatic events was measured using the Harvard Trauma Questionnaire-part I, while psychological distress and depressive symptoms were assessed with the General Health Questionnaire and the Patient Health Questionnaire. After three and six months of follow-up, the proportion of participants who developed a mental disorder was calculated using the Mini International Neuropsychiatric Interview.

Results: A total of 1,101 participants were included in the analysis. At three- and six-month follow-up the observed frequency of mental disorders was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. After controlling for confounders, logistic regression analysis showed that participants with a lower education level ($p = 0.034$), a shorter duration of journey ($p = 0.057$), and arriving from countries with war-related contexts ($p = 0.017$), were more at risk of developing mental disorders. Psychological distress ($p = 0.004$), depression ($p = 0.001$), and

1
2
3
4 exposure to potentially traumatic events ($p = 0.020$) were predictors of mental disorder
5
6 development.
7
8

9 **Conclusions:** This study identified several risk factors for the development of mental disorders
10
11 in asylum seekers and refugees, some of which may be the target of risk reduction policies. The
12
13 identification of asylum seekers and refugees at increased risk of mental disorders should guide
14
15 the implementation of focused preventative psychological interventions.
16
17
18
19

20
21
22 **Key-words:** asylum seekers, refugees, migrants, mental health, mental disorders, high-income
23
24 countries, low-income countries
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

According to the last United Nations Refugee Agency data, despite COVID-related movement restrictions, at the end of 2020 more than 82 million people were forcibly displaced as a result of persecution, conflict, violence, and other human rights violations (UNHCR, 2022). Of these, more than 30 million were asylum seekers or refugees. While Turkey continued to host the largest number of refugees, most of whom were Syrian refugees, 10% of all the world's refugees were resettled in Europe at the end of 2020 (UNHCR, 2022). These numbers are expected to further increase, as recent estimates have suggested that over 6.5 million refugees from Ukraine have crossed to Poland, Hungary, Romania, Moldova and other countries (UNHCR, 2022).

Despite heterogeneity between the studies and the populations of refugees and asylum seekers assessed (Giacco, 2019), a large body of evidence consistently showed that the frequency of mental disorders in refugees and asylum seekers is increased as compared with the general population. Existing World Health Organization (WHO) estimates suggest a prevalence of 13.0% for mild forms of depression, anxiety, and post-traumatic stress disorder (PTSD), and 4.0% for moderate forms (Charlson *et al.*, 2019). Other reviews of prevalence studies found adult refugee and asylum seekers have high and persistent rates of depression and PTSD, while the prevalence of anxiety disorders and psychosis are more comparable to findings from general populations (Blackmore *et al.*, 2020, Henkelmann *et al.*, 2020).

The increased frequency of mental disorders in refugees and asylum seekers may be related to the refugee experience, which is characterized by loss of homes, hopes, possessions, and disruption of personal, family, and professional life projects. In addition, a wide range of physical, psychological, and psychosocial problems associated with adversities may occur, such as bombings, threats, captivity, torture, injury, witnessing death or injury of loved ones, discrimination, economic stress, and uncertainty about the future (Priebe *et al.*, 2016).

1
2
3
4 Interestingly, while these factors have been studied as determinants of poor mental health
5 outcomes, or as determinants of psychological distress (Jannesari *et al.*, 2020), to our knowledge
6 they have never been investigated in this population as risk factors for the development of
7 mental disorders considered as full-blown diagnostic entities (Gleeson *et al.*, 2020, Hajak *et al.*,
8 2021). Knowledge of determinants of the development of mental disorders is particularly
9 important to implement policies aiming at decreasing exposure to such determinants, to
10 improve identification of refugees and asylum seekers at increased risk of developing mental
11 disorders, and to plan the provision of psychological interventions to individuals and
12 communities at risk.

13
14
15
16
17
18
19
20
21
22
23
24
25 Aiming to test the effect of Self Help Plus (SH+), a low intensity, group-based, self-help
26 psychological intervention recently developed by the WHO (Epping-Jordan *et al.*, 2016, World
27 Health Organization, 2021a), in reducing the frequency of mental disorders, we conducted two
28 large randomized prevention trials in asylum seekers and refugees, one in western Europe and
29 another in Turkey (Purgato *et al.*, 2019). The studies followed the same research protocol and
30 were conducted in parallel (Acarturk *et al.*, 2022, Purgato *et al.*, 2021). Both the Western
31 European and Turkey studies showed evidence of an effect of SH+ in preventing the onset of
32 mental disorders, but differences were observed between the studies. The effect was much
33 more pronounced for the Turkey study where efficacy (i.e. reducing the frequency of any mental
34 disorder) was observed at 6 months, compared to the Western European study where a
35 preventative effect was only found post-intervention and not after six months (Acarturk *et al.*,
36 2022, Purgato *et al.*, 2021). As these are the only two studies that enrolled asylum seekers and
37 refugees without any mental disorders at baseline, and that assessed the frequency of mental
38 disorders at follow-up as primary outcome (Papola *et al.*, 2020), they offered a unique
39 opportunity to prospectively investigate the risk factors for the onset of mental disorders in a
40 large group of asylum seekers and refugees.

Methods

Participants and measures

Participant-level data from two randomized prevention trials involving asylum seekers and refugees resettled in Western European countries (Austria, Finland, Germany, Italy, and UK) and in Turkey were pooled (Acarturk *et al.*, 2022, Purgato *et al.*, 2021). In both studies, participants were randomly assigned to the SH+ psychological intervention, consisting of SH+ combined with enhanced care as usual (ECAU), or to ECAU only. The two studies were conducted in parallel following the same study design (Purgato *et al.*, 2019). The protocol of the present study was registered within the Open Science Framework (<https://osf.io/37h5n>).

In both studies, participants were included if they met the following criteria: a) aged 18 years or older; b) able to speak and understand Arabic, Dari, English, or Urdu; c) being under temporary protection with a refugee or asylum seeker status; d) experiencing psychological distress, as shown by a score of 3 or more on the 12-item dichotomously-scored General Health Questionnaire (el-Rufaie and Daradkeh, 1996, Goldberg *et al.*, 1998, Kilic *et al.*, 1997); e) having completed oral and written informed consent to enter the study. Exclusion criteria were: a) presence of any mental disorder according to the Mini International Neuropsychiatric Interview (M.I.N.I.), a brief structured diagnostic interview for the major psychiatric disorders in DSM-III-R, DSM-IV and DSM-5 and ICD-10. (Kadri *et al.*, 2005, Sheehan *et al.*, 1998); b) evidence of acute medical conditions contraindicating study participation; c) evidence of imminent suicide risk, or suicide risk scored as “moderate or high” on the M.I.N.I.; d) signs of impaired decision-making capacity emerging from responses during the clinical interview.

The M.I.N.I. was administered before random allocation, in order to exclude participants with a mental disorder, and at three and six months of follow-up, in order to calculate the proportion

1
2
3
4 of participants who developed a mental disorder. Exposure to potentially traumatic events was
5
6 measured at baseline using the Harvard Trauma Questionnaire-part I (HTQ) (Mollica *et al.*,
7
8 1992). The HTQ-part I covers a variety of trauma events that may affect refugee mental health,
9
10 and the scoring represents the number of different types of traumatic events experienced by
11
12 the participants (higher score is associated with high number of traumatic events). In addition
13
14 to using the HTQ total score, we identified subtypes of traumatic events performing a Principal
15
16 Components analysis (PCA).
17
18

19
20
21 Mental health symptoms were measured using the following validated instruments. The GHQ-
22
23 12 questionnaire was used to measure psychological distress (el-Rufaie and Daradkeh, 1996,
24
25 Goldberg *et al.*, 1998, Kilic *et al.*, 1997), while PTSD was assessed with the PTSD Checklist for
26
27 DSM-5 (PCL-5), a 20-item questionnaire that measures overall PTSD symptoms (score zero to
28
29 80) and symptoms by cluster (intrusions, avoidance, negative changes in thoughts and mood,
30
31 and changes in arousal), with higher scores indicating higher levels of PTSD symptoms
32
33 (Blanchard *et al.*, 1996). Levels of self-reported depression symptoms were measured with the
34
35 Patient Health Questionnaire, nine-item version (PHQ-9) which has a four-point scale (score 0
36
37 to 27) (Kroenke *et al.*, 2001). All measures were collected at baseline before random allocation,
38
39 and after three and six months of follow-up. Assessors were trained in the administration of
40
41 rating scales, instructed on how to perform follow-up assessments while preserving effective
42
43 masking, and assisted by cultural mediators when needed.
44
45
46
47

48 *Interventions*

49
50
51 The SH+ intervention was developed by the WHO, as described elsewhere (Acarturk *et al.*, 2022,
52
53 Purgato *et al.*, 2021), and is now publicly available (World Health Organization, 2021a). SH+
54
55 consists of a pre-recorded audio course complemented with an illustrated self-help book. The
56
57 book has been recently updated and published by WHO as *Doing What Matters in Times of*
58
59
60

1
2
3
4 Stress (World Health Organization, 2021b). All SH+ materials were adapted for the cultural
5
6 groups included in the studies. The SH+ pre-recorded audio material was delivered across five
7
8 2-hour sessions to groups of up to 30 people. The audio material imparts key information about
9
10 stress management and guides participants through individual exercises and small group
11
12 discussions. To augment the audio recordings, an illustrated self-help book reviews all essential
13
14 content and concepts. The SH+ intervention was fully delivered in the language of participants
15
16 by trained facilitators with a migration background, who were native speakers of the target
17
18 languages. ECAU, provided both to the experimental and control group, consisted of routinely
19
20 delivered social support and/or care according to local regulations. Participants in the ECAU arm
21
22 received the same baseline and follow-up assessments of the intervention arm, according to the
23
24 study schedule (around three and six months after randomization), information about freely
25
26 available health and social services, and links to community networks providing support to
27
28 refugees and asylum seekers.
29
30
31
32

33 34 *Statistical analysis*

35
36 Descriptive statistics (mean and SD for continuous variables and absolute numbers and
37
38 percentages for dichotomous variables) were computed on sociodemographic, premigration,
39
40 migration and postmigration variables at baseline, and for clinical variables.
41
42
43

44 Participants who met criteria for any mental disorder on the M.I.N.I. at three or six-month
45
46 follow-up were considered cases with a mental disorder. Multiple imputation was adopted to
47
48 address the issue of missing data in all the variables included in the model. In particular, in case
49
50 of missing data at the M.I.N.I. in one timepoint, imputation was performed on single M.I.N.I.
51
52 values at three and six months. In case of missing values on continuous clinical measures, the
53
54 imputation was performed on single item scales. Specifically, imputation followed the approach
55
56 reported by Plumpton and colleagues (Plumpton *et al.*, 2016), that is we used scale totals within
57
58
59
60

1
2
3
4 prediction equations and, for imputations of responses to individual scale items, we additionally
5
6 included the responses to the other scale items, using the “ice” Stata routine (Royston, 2005,
7
8 Sterne *et al.*, 2009), and considering single-item scores as ordered categorical variables. M.I.N.I.
9
10 values at the two time-points were used in the prediction equations of regression predictors,
11
12 and upper and lower bounds were set for continuous variables with missing values as
13
14 appropriate. The number of imputed samples was determined by following the rule of thumb
15
16 suggested by White and colleagues, i.e.: “at least equal to the percentage of incomplete cases”
17
18 (White *et al.*, 2011). We rounded such number to the nearest multiple of 10 above.
19
20

21
22
23 In order to identify subtypes of traumatic events, we performed a PCA on tetrachoric
24
25 correlations of HTQ items, with Quartimin oblique rotation to allow for between-factor non-null
26
27 correlations (Tabachnick, 2007), and summing items with loadings above 0.40 on the same
28
29 factor to create scores for the regression model (Stevens, 2002). The number of factors was
30
31 determined by adopting Kaiser's rule (Kaiser, 1960), i.e. using the cut-off scores of 1 for the
32
33 eigenvalues.
34
35

36
37 In order to investigate predictors of the development of mental disorders, unadjusted and
38
39 adjusted logistic regression analyses were performed, using the frequency of participants with
40
41 a mental disorder in at least one timepoint as binary dependent variable. The following
42
43 independent variables were inserted into the model: age (years), gender (men, women),
44
45 education (years), unemployment (yes/no), country of origin (Syria, Iraq, Nigeria, other
46
47 countries), length of journey (below one month, between one and three months, above three
48
49 months), study (Western Europe versus Turkey), length of stay in the resettlement country
50
51 (months), HTQ (total score at baseline), GHQ-12 (total score at baseline), PCL-5 (total score at
52
53 baseline), PHQ-9 (total score at baseline), number of SH+ sessions received (zero to five).
54
55 Considering that PTSD and depression are known to be interrelated experiences following
56
57
58
59
60

1
2
3
4 trauma (Rytwinski *et al.*, 2013), and considering that the PCL-5 largely overlaps with the HTQ
5
6 (Patel *et al.*, 2022), and that a substantial overlap exists between the PCL-5 and the PHQ-9
7
8 (Dabrowski CL, 2020), we excluded the PCL-5 from the final model, but it was included (total
9
10 score at baseline) in the imputation model.
11
12

13
14 In order to check the robustness of the analysis, we re-ran the model after excluding the arm
15
16 receiving SH+. As a further sensitivity analysis, in order to estimate within-centre effects, we
17
18 performed a model including a fixed effect for recruiting centre. As a subgroup analysis, we re-
19
20 ran the logistic regression analysis separately for each study sample (western Europe versus
21
22 Turkey). All analyses were performed using Stata 17 (Statacorp, 2017).
23
24
25
26
27

28 **Results**

29 *Participants*

30
31
32
33
34 The two prevention trials randomized a total of 1,101 participants. After six months, 249
35
36 participants (22.6%) were not available for follow-up assessments, for the reasons reported in
37
38 Figure 1. Participant characteristics at baseline are presented in Table 1. Almost half were
39
40 female, with a mean age of 32 (SD 9.521) years, and a mean education of 10 years (SD 4.424).
41
42 Slightly less than 40% was unemployed. The migration journey lasted more than three months
43
44 in around one third of participants, in another one third it lasted between one and three months,
45
46 while in the remaining 40% the host country was reached in less than one month. On average,
47
48 the mean length of stay in the resettlement country was slightly more than three years. Most of
49
50 the participants were from Syria, Nigeria and Iraq (Table 1). Participants attended a mean
51
52 number of 1.4 (SD 2.015) SH+ sessions. The mean (SD) baseline scores on the measures of
53
54 interest are reported in Table 1.
55
56
57
58
59
60

Frequency and determinants of mental disorders

At three- and six-month follow-up the observed frequency of mental disorders, as measured with the M.I.N.I., was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. The majority of detected mental disorders were major depressive disorders (9.5% and 20.3% after three and six months), PTSD (3.2% and 6.7%), and anxiety disorders (2.1% and 4.1%) (Figure 1).

The PCA model found a 3-factor solution where the set of items was exhaustive and mutually exclusive, with the HTQ items grouped as follows: factor 1 - lack of basic needs: lack of food or water, ill health without access to medical care, lack of shelter; factor 2 - violence and abuse: imprisonment, serious injury, brain washing, rape or sexual abuse, forced isolation from others, forced separation from family members, torture, other (e.g. domestic violence;) factor 3 - being close to death: combat situation, being close to death, murder of family or friend, unnatural death of family or friend, murder of stranger(s), lost or kidnapped. The factor loadings of the PCA-model with Quartimin oblique rotation are reported in the Supplemental Materials.

The results of unadjusted and adjusted logistic regression analyses investigating factors associated with the development of mental disorders are presented in Table 2. Participants with lower education level ($p = 0.034$), a shorter duration of journey ($p = 0.057$), and arriving from countries with war-related contexts (Iraq, Syria) ($p = 0.017$), were those more at risk of developing mental disorders. In terms of mental health symptoms and exposure to traumatic events, psychological distress ($p = 0.004$), depressive symptoms ($p = 0.001$), and HTQ total score ($p = 0.020$) at baseline were predictors of mental disorder development, as was the HTQ factor violence and abuse (imprisonment, serious injury, brain washing, rape or sexual abuse, forced isolation from others, forced separation from family members, torture, domestic violence) ($p =$

1
2
3
4 0.008). In terms of post-migration factors, the number of SH+ sessions ($p < 0.001$) was inversely
5
6 associated with the risk of developing mental disorders (Table 2).
7

8
9 Secondary logistic regression analyses including only the ECAU arms of the two studies, and
10
11 analysing the two studies separately, confirmed the role of mental health symptoms and
12
13 exposure to traumatic events as risk factors for mental disorders (Supplemental Materials).
14
15 However, due to lower statistical power, some factors lost significance despite a similar or even
16
17 higher estimated effect, as for example the HTQ in the model restricted to the western European
18
19 study, or years of education and travel duration in the model restricted to the ECAU sample, and
20
21 GHQ-12 in the model restricted to the study conducted in Turkey (Supplemental Materials).
22
23

24
25 Consistently with the results of the two studies, the number of SH+ sessions was inversely
26
27 associated with the frequency of mental disorders only in the study conducted in Turkey
28
29 (Supplemental Materials). The inclusion of recruiting centres as predictor variables revealed that,
30
31 in comparison with participants recruited in Istanbul, participants recruited in Vienna were less
32
33 likely to develop a mental disorder (Supplemental Materials).
34
35
36
37
38
39

40 **Discussion**

41
42
43 To the best of our knowledge, this is the first prospective study that examined risk factors for
44
45 the onset of mental disorders among asylum seekers and refugees without a mental disorder at
46
47 baseline. We showed the significant influence of socio-demographic, clinical, and contextual
48
49 factors, including potentially traumatic events, on the development of mental disorders in
50
51 asylum seekers and refugees resettled in Western European countries and in Turkey.
52
53

54
55 Overall, one in four refugees and asylum seekers developed a diagnosable mental disorder over
56
57 a period of six months. Interestingly, the most frequently reported diagnostic group was
58
59
60

1
2
3
4 depression, followed by PTSD and anxiety disorders. These figures, derived from two
5
6 intervention studies, cannot be compared with those from epidemiological studies, which
7
8 generally show similar frequencies for depression and PTSD, or slightly higher frequencies for
9
10 depression, depending on the population surveyed and the study setting (Blackmore *et al.*, 2020,
11
12 Charlson *et al.*, 2019, Henkelmann *et al.*, 2020, Hoell *et al.*, 2021). In migrants exposed to armed
13
14 conflict, by contrast, the frequency of PTSD was found to be higher than depression (31% vs
15
16 25%) (Mesa-Vieira *et al.*, 2022).
17
18
19

20
21 The finding that educational level is inversely associated with the development of mental
22
23 disorders expands previous data collected in the general population showing that higher
24
25 educational level seems to have a protective effect against anxiety and depression symptoms,
26
27 and against common mental disorders in general (Araya *et al.*, 2003, Bjelland *et al.*, 2008). In the
28
29 general population, poor education has been suggested to be a marker of lack of opportunities
30
31 and resources, including material and psychological resources, and a marker of childhood
32
33 adversity (Araya *et al.*, 2003). In asylum seekers and refugees, these factors may directly
34
35 contribute to the emergence of mental disorders, or may interfere with the coping skills required
36
37 to deal with all the adversities associated with the migration and resettlement process, which,
38
39 in turn, may increase the risk of developing mental disorders (Kirmayer *et al.*, 2011).
40
41
42

43
44 Consistently with an extensive literature showing that exposure to potentially traumatic events
45
46 represents a risk factor for poor mental health and well-being (Gleeson *et al.*, 2020, Hajak *et al.*,
47
48 2021, Priebe *et al.*, 2016), the present study adds that exposure to potentially traumatic events
49
50 is also a risk factor for the development of mental disorders considered as full-blown diagnostic
51
52 entities. The finding that being displaced from countries with war-related contexts emerged as
53
54 an additional risk factor for mental disorders further corroborates this association. War-related
55
56 contexts imply exposure to multiple and serious traumatic events, and the exposure to combat
57
58
59
60

1
2
3
4 situations creates the risk of witnessing violence and/or death and experiencing physical and
5
6 psychological violence (Crepet *et al.*, 2017). Extensive literature shows that these situations are
7
8 associated with poor mental health outcomes, including PTSD, depression, anxiety, and
9
10 somatization symptoms (Jongedijk *et al.*, 2020, Knipscheer *et al.*, 2015, Nickerson *et al.*, 2021).

11
12
13
14 The present study has limitations and strengths. A first limitation is that the population enrolled
15
16 in the two trials cannot be considered representative of the general population of asylum
17
18 seekers and refugees, as we selected participants scoring above a threshold of psychological
19
20 distress, and we excluded those at suicide risk. During the studies, in addition, around half of
21
22 the participants was exposed to a psychological intervention aimed at preventing the
23
24 development of mental disorders. For these reasons, the overall frequency estimate of mental
25
26 disorders cannot apply to the general population of asylum seekers and refugees resettled in
27
28 Western European countries or in Turkey. A second issue is that we did not calculate a true
29
30 incidence of mental disorders, but only frequency figures at two time points with a structured
31
32 diagnostic interview that might overestimate the true frequency of some mental disorders, such
33
34 as depression (Wu *et al.*, 2020). Third, the SARS-CoV2 pandemic impacted the study procedures,
35
36 because in all the recruiting sites follow-up assessments were conducted using online tools
37
38 instead of face-to-face meetings. Although assessors were trained in the administration of rating
39
40 scales, instructed on how to perform online follow-up assessments, and assisted by cultural
41
42 mediators when needed, it is unknown whether this may have impacted the responses of
43
44 participants to the instruments. Related to this, even though several studies documented that a
45
46 careful and culturally appropriate use of available instruments is feasible and allows a
47
48 standardization of the screening process and a systematic recognition of psychological distress
49
50 and psychiatric diagnoses (Acarturk *et al.*, 2021) we acknowledge that formal studies on use of
51
52 these tools in refugee groups are lacking. Fourth, despite a growing body of literature showed
53
54 that the duration of the asylum procedure is an important risk factor for mental health
55
56
57
58
59
60

1
2
3
4 conditions (Laban et al. 2004; Winkler et al. 2019), we acknowledge that this information was
5
6 not collected. However, months in host country, which may be considered a proxy of length of
7
8 asylum procedure, was included in the model but did not emerge as a significant factor.
9

10
11 Despite these limitations, there are strengths that should be emphasised. The main strength is
12
13 the exclusion of participants with a mental disorder at baseline, and the choice of a dependent
14
15 variable that is fully consistent with a preventative design, namely the frequency of mental
16
17 disorders at follow-up, assessed with validated measures. This design allowed to prospectively
18
19 investigate the determinants of mental disorders rather than of poor mental health, as previous
20
21 studies have done in populations who might already be with mental disorders at baseline (Priebe
22
23 *et al.*, 2016). Another strength is a sample size of over a thousand of asylum seekers and
24
25 refugees, resettled in a variety of western European sites and in Turkey. This aspect is of
26
27 relevance not only in terms of statistical power, but also in terms of generalisability and
28
29 applicability of study findings to different types of reception settings in high-income and middle-
30
31 income countries. Lastly, despite an attrition rate of around 20%, a follow-up assessment of six
32
33 months is noteworthy in such a difficult-to-follow population, who is often moved from one
34
35 reception site to another, and may not perceive mental health as a priority, having a number of
36
37 other challenging concerns such as housing, unstable working conditions, management of visa
38
39 issues, safety of family members, fear of being returned to home country, plans to move to
40
41 another country or to another location.
42
43
44
45
46
47

48 The present study has important practical implications. The finding of a positive relationship
49
50 between exposure to potentially traumatic events and risk of mental disorder development
51
52 suggests a pressing need of developing policies aiming to decrease exposure to such traumatic
53
54 experiences after resettlement. Host countries may have opportunities to decrease exposure to
55
56 situations such as material and economic hardship that could affect integrity, independence,
57
58
59
60

1
2
3
4 dignity and well-being (financial strain), social hardship due to loss of status (social strain),
5
6 feelings of inadequacy in relation with specific skills needed in the host-country to successfully
7
8 function in daily life (competency strain), experiences of unfair treatment on the basis of
9
10 prejudice (perceived discrimination). This ambitious goal may be achieved by implementing
11
12 reception conditions that optimize internationally recognized minimal quality standards. For
13
14 example, standards for the reception of applicants for international protection have been
15
16 established by Directive 2013/33/EU of the European Parliament. The Directive clearly reports
17
18 that national authorities should ensure that reception modalities are specifically designed to
19
20 meet the needs of persons requiring international protection, including legal assistance,
21
22 document provision, material support, links with local communities, freedom of movement,
23
24 information about labor market access, vocational training, social support. Health care, including
25
26 mental health care, is also mentioned as a key intervention where needed. It would be important
27
28 to ensure that efforts to support forcibly displaced people are coordinated across the different
29
30 layers of the social environments in which they are hosted, i.e. at the level of the individual, their
31
32 family, the community, and the institutions that have governance responsibility for their care
33
34 and support (White and Van der Boor, 2021).
35
36
37
38
39
40

41 In addition to implications in terms of risk reduction policies, the present study has implications
42
43 in terms of provision of psychological interventions aimed at preventing the development of
44
45 mental disorders (Miller *et al.*, 2021). As the implementation of preventative psychological
46
47 interventions to the whole population of asylum seekers and refugees may not be sustainable
48
49 by host countries, national authorities may consider to offer psychological interventions to at-
50
51 risk populations. The finding that persons with lower education level, a shorter duration of
52
53 journey, arriving from countries with war-related contexts (Iraq, Syria), and with high level of
54
55 psychological distress and depressive symptoms, are those more at risk of developing mental
56
57 disorders, appears to identify a target population for focused psychological prevention
58
59
60

1
2
3
4 interventions. The WHO has recently developed a number of low-intensity psychological
5
6 interventions that may be scaled up as public health strategies to address mental disorders and
7
8 psychological distress in refugee populations exposed to ongoing adversities (World Health
9
10 Organization, 2017). In addition to WHO interventions, other psychological treatments have
11
12 been shown to be effective in alleviating psychological symptoms in asylum seekers and
13
14 refugees (Turrini *et al.*, 2021, Uphoff *et al.*, 2020), but a preventative effect has been
15
16 investigated for SH+ only. The present results, by showing an inverse association between
17
18 number of SH+ sessions and risk of developing a mental disorder, further corroborate the value
19
20 of SH+ as a prevention intervention. The finding that trauma exposure stands out as a predictor
21
22 for development of mental disorders would additionally suggest that interventions may need to
23
24 include a focus on traumatic memories and experiences, in addition to being focused on low
25
26 mood.
27
28
29
30

31
32 In parallel with the provision of psychological interventions to at-risk populations, national
33
34 authorities should ensure regular access to such interventions supporting at risk populations to
35
36 engage through reducing barriers that might prevent or limit access or use (Fuhr *et al.*, 2019).
37
38 This can be achieved by optimizing access (e.g. accessible services and supports) to a range of
39
40 interventions depending on need (e.g. stepped care), and by using different delivery
41
42 mechanisms (e.g. digital or peer delivered support).
43
44
45

46
47 As the number of persons in need of protection is likely to substantially increase globally, driven
48
49 by long-lasting wars as well as by new conflicts such as the Russo-Ukrainian war which broke out
50
51 recently (Barbui *et al.*, 2022, Pandi-Perumal *et al.*, 2022), national authorities are urged to
52
53 develop reception and resettlement programs meeting the needs of this vulnerable group.
54
55 These programs should be designed and implemented attempting to decrease the risk of post-
56
57 migration stressors that may contribute to worsening the mental health of a population already
58
59
60

1
2
3
4 exposed to potentially traumatic experiences before and during the migration process.
5
6 Evidence-based focused psychological support should be an important program component to
7
8 be delivered to asylum seekers and refugees at-risk of developing a mental disorder.
9
10
11
12

13 **Contributors** CB, MP, and FT conceived the study. FT statistically analysed the data. CB, GO
14 and FT accessed and verified the data. CB wrote the first draft of the manuscript with input
15 from FT and MP. All authors contributed to the conception and design of the study, data
16 interpretation and manuscript revision. All authors read and approved the submitted
17 manuscript and had final responsibility for the decision to submit for publication. CB was the
18 overall guarantor of the content.
19
20
21

22 **Funding** This work was supported by the European Commission (grant agreement No. 779255;
23 “RE-DEFINE: Refugee Emergency: DEFining and Implementing Novel Evidence-based
24 psychosocial interventions”).
25
26
27

28 **Disclaimer** The authors alone are responsible for the views expressed in this article, and they
29 do not necessarily represent the views, decisions, or policies of the institutions with which they
30 are affiliated.
31
32
33

34 **Competing interests** None declared.
35
36
37

38 **Ethics approval** This was a secondary analysis of deidentified participant data. The initial study
39 was approved by the WHO Research Ethics Review Committee, and from the Ethics
40 Committees of all participating sites. Before participation, a written informed consent was
41 obtained from each participant, allowing investigators to use anonymised data for future
42 analysis. Participants gave informed consent to participate in the study before taking part.
43
44
45

46 **Data availability statement** Data collected in the RE-DEFINE study are stored in the online
47 repository EUDAT B2SHARE (study in Western Europe:
48 <https://b2share.eudat.eu/records/fa7264d624364683830ff37acee01c04>; study in Turkey:
49 <http://doi.org/10.23728/b2share.8ac4f28d2415413e89de7847c05471fc>). The reuse of data
50 will be offered only upon motivated request, which will undergo the scrutiny of the RE-DEFINE
51 General Assembly.
52
53
54
55
56
57
58
59
60

References

- Acarturk, C., McGrath, M., Roberts, B., Ilkkursun, Z., Cuijpers, P., Sijbrandij, M., Sondorp, E., Ventevogel, P., McKee, M. & Fuhr DC. (2021) Prevalence and predictors of common mental disorders among Syrian refugees in Istanbul, Turkey: a cross-sectional study. *Social Psychiatry and Psychiatric Epidemiology* 56, 475-84.
- Acarturk, C., Uygun, E., Ilkkursun, Z., Carswell, K., Tedeschi, F., Batu, M., Eskici, S., Kurt, G., Anttila, M., Au, T., Baumgartner, J., Churchill, R., Cuijpers, P., Becker, T., Koesters, M., Lantta, T., Nose, M., Ostuzzi, G., Popa, M., Purgato, M., Sijbrandij, M., Turrini, G., Valimaki, M., Walker, L., Wancata, J., Zanini, E., White, R. G., van Ommeren, M. & Barbui, C. (2022). Effectiveness of a WHO self-help psychological intervention for preventing mental disorders among Syrian refugees in Turkey: a randomized controlled trial. *World Psychiatry* 21, 88-95.
- Araya, R., Lewis, G., Rojas, G. & Fritsch, R. (2003). Education and income: which is more important for mental health? *Journal of Epidemiology and Community Health* 57, 501-5.
- Barbui, C., Purgato, M., Acarturk, C., Churchill, R., Cuijpers, P., Koesters, M., Sijbrandij, M., Valimaki, M., Wancata, J. & White, R. G. (2022). Preventing the mental health consequences of war in refugee populations. *Epidemiology and Psychiatric Sciences* 31, e24.
- Bjelland, I., Krokstad, S., Mykletun, A., Dahl, A. A., Tell, G. S. & Tambs, K. (2008). Does a higher educational level protect against anxiety and depression? The HUNT study. *Social Science and Medicine* 66, 1334-45.
- Blackmore, R., Boyle, J. A., Fazel, M., Ranasinha, S., Gray, K. M., Fitzgerald, G., Misso, M. & Gibson-Helm, M. (2020). The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis. *PLoS Medicine* 17, e1003337.
- Blanchard, E. B., Jones-Alexander, J., Buckley, T. C. & Forneris, C. A. (1996). Psychometric properties of the PTSD Checklist (PCL). *Behavioural Research and Therapeutics* 34, 669-73.
- Charlson, F., van Ommeren, M., Flaxman, A., Cornett, J., Whiteford, H. & Saxena, S. (2019). New WHO prevalence estimates of mental disorders in conflict settings: a systematic review and meta-analysis. *Lancet* 394, 240-248.
- Crepet, A., Rita, F., Reid, A., Van den Boogaard, W., Deiana, P., Quaranta, G., Barbieri, A., Bongiorno, F. & Di Carlo, S. (2017). Mental health and trauma in asylum seekers landing in Sicily in 2015: a descriptive study of neglected invisible wounds. *Conflict and Health* 11, 1.
- Dabrowski CL, L. J., Sherman MF, Kirkhart MW, Breen D (2020). Probable Ptsd, Symptoms of Depression, and an Exploratory Factor Analysis of Item Responses in Canadian Steelworkers After Workplace Traumas. *Crisis, Stress, and Human Resilience: An International Journal* 2.

- 1
2
3
4 el-Rufaie, O. F. & Daradkeh, T. K. (1996). Validation of the Arabic versions of the thirty- and
5 twelve-item General Health Questionnaires in primary care patients. *British Journal of*
6 *Psychiatry* 169, 662-4.
7
8
9 Epping-Jordan, J. E., Harris, R., Brown, F. L., Carswell, K., Foley, C., Garcia-Moreno, C., Kogan, C.
10 & van Ommeren, M. (2016). Self-Help Plus (SH+): a new WHO stress management
11 package. *World Psychiatry* 15, 295-296.
12
13 Fuhr, D. C., Acarturk, C., McGrath, M., Ilkkursun, Z., Sondorp, E., Sijbrandij, M., Ventevogel, P.,
14 Cuijpers, P., McKee, M. & Roberts, B. (2019). Treatment gap and mental health service
15 use among Syrian refugees in Sultanbeyli, Istanbul: a cross-sectional survey. *Epidemiology*
16 *and Psychiatric Sciences* 29, e70.
17
18 Giacco, D. (2019). Identifying the critical time points for mental health of asylum seekers and
19 refugees in high-income countries. *Epidemiology and Psychiatric Sciences* 29, e61.
20
21 Gleeson, C., Frost, R., Sherwood, L., Shevlin, M., Hyland, P., Halpin, R., Murphy, J. & Silove, D.
22 (2020). Post-migration factors and mental health outcomes in asylum-seeking and
23 refugee populations: a systematic review. *European Journal of Psychotraumatology* 11,
24 1793567.
25
26 Goldberg, D. P., Oldehinkel, T. & Ormel, J. (1998). Why GHQ threshold varies from one place to
27 another. *Psychological Medicine* 28, 915-21.
28
29 Hajak, V. L., Sardana, S., Verdeli, H. & Grimm, S. (2021). A Systematic Review of Factors Affecting
30 Mental Health and Well-Being of Asylum Seekers and Refugees in Germany. *Frontiers in*
31 *Psychiatry* 12, 643704.
32
33 Henkelmann, J. R., de Best, S., Deckers, C., Jensen, K., Shahab, M., Elzinga, B. & Molendijk, M.
34 (2020). Anxiety, depression and post-traumatic stress disorder in refugees resettling in
35 high-income countries: systematic review and meta-analysis. *BJPsych Open* 6, e68.
36
37 Hoell, A., Kourmpeli, E., Salize, H. J., Heinz, A., Padberg, F., Habel, U., Kamp-Becker, I., Hohne, E.,
38 Boge, K. & Bajbouj, M. (2021). Prevalence of depressive symptoms and symptoms of post-
39 traumatic stress disorder among newly arrived refugees and asylum seekers in Germany:
40 systematic review and meta-analysis. *BJPsych Open* 7, e93.
41
42 Jannesari, S., Hatch, S. & Oram, S. (2020). Seeking sanctuary: rethinking asylum and mental
43 health. *Epidemiology and Psychiatric Sciences* 29, e154.
44
45 Jongedijk, R. A., Eising, D. D., van der Aa, N., Kleber, R. J. & Boelen, P. A. (2020). Severity profiles
46 of posttraumatic stress, depression, anxiety, and somatization symptoms in treatment
47 seeking traumatized refugees. *Journal of Affective Disorders* 266, 71-81.
48
49 Kadri, N., Agoub, M., El Gnaoui, S., Alami Kh, M., Hergueta, T. & Moussaoui, D. (2005). Moroccan
50 colloquial Arabic version of the Mini International Neuropsychiatric Interview (MINI):
51 qualitative and quantitative validation. *European Psychiatry* 20, 193-5.
52
53 Kaiser, H. F. (1960). The Application of Electronic Computers to Factor Analysis. *Educational and*
54 *Psychological Measurement* 20.
55
56
57
58
59
60

- 1
2
3
4 Kilic, C., Rezaki, M., Rezaki, B., Kaplan, I., Ozgen, G., Sagduyu, A. & Ozturk, M. O. (1997). General
5 Health Questionnaire (GHQ12 & GHQ28): psychometric properties and factor structure of
6 the scales in a Turkish primary care sample. *Social Psychiatry and Psychiatric Epidemiology*
7 32, 327-31.
8
9
10 Kirmayer, L. J., Narasiah, L., Munoz, M., Rashid, M., Ryder, A. G., Guzder, J., Hassan, G., Rousseau,
11 C., Pottie, K., Canadian Collaboration for, I. & Refugee, H. (2011). Common mental health
12 problems in immigrants and refugees: general approach in primary care. *CMAJ* 183, E959-
13 67.
14
15 Knipscheer, J. W., Sleijpen, M., Mooren, T., Ter Heide, F. J. & van der Aa, N. (2015). Trauma
16 exposure and refugee status as predictors of mental health outcomes in treatment-
17 seeking refugees. *BJPsych Bull* 39, 178-82.
18
19
20 Kroenke, K., Spitzer, R. L. & Williams, J. B. (2001). The PHQ-9: validity of a brief depression
21 severity measure. *Journal of General Internal Medicine* 16, 606-13.
22
23 Laban, C.J., Gernaat, H.B., Komproe, I.H., Schreuders, B.A. & De Jong, J.T. (2004). Impact of a
24 long asylum procedure on the prevalence of psychiatric disorders in Iraqi asylum seekers
25 in The Netherlands. *Journal of Nervous and Mental Disorders* 192, 843-51.
26
27
28 Mesa-Vieira, C., Haas, A. D., Buitrago-Garcia, D., Roa-Diaz, Z. M., Minder, B., Gamba, M.,
29 Salvador, D., Jr., Gomez, D., Lewis, M., Gonzalez-Jaramillo, W. C., Pahud de Mortanges, A.,
30 Buttia, C., Muka, T., Trujillo, N. & Franco, O. H. (2022). Mental health of migrants with pre-
31 migration exposure to armed conflict: a systematic review and meta-analysis. *Lancet*
32 *Public Health* 7, e469-e481.
33
34
35 Miller, K. E., Jordans, M. J. D., Tol, W. A. & Galappatti, A. (2021). A call for greater conceptual
36 clarity in the field of mental health and psychosocial support in humanitarian settings.
37 *Epidemiology and Psychiatric Sciences* 30, e5.
38
39
40 Mollica, R. F., Caspi-Yavin, Y., Bollini, P., Truong, T., Tor, S. & Lavelle, J. (1992). The Harvard
41 Trauma Questionnaire. Validating a cross-cultural instrument for measuring torture,
42 trauma, and posttraumatic stress disorder in Indochinese refugees. *Journal of Nervous*
43 *and Mental Disorders* 180, 111-6.
44
45
46 Nickerson, A., Byrow, Y., Rasmussen, A., O'Donnell, M., Bryant, R., Murphy, S., Mau, V.,
47 McMahon, T., Benson, G. & Liddell, B. (2021). Profiles of exposure to potentially traumatic
48 events in refugees living in Australia. *Epidemiology and Psychiatric Sciences* 30, e18.
49
50
51 Pandi-Perumal, S. R., Kumar, V. M., Pandian, N. G., de Jong, J. T., Andiappan, S., Corlățeanu, A.,
52 Mahalaksmi, A. M., Chidambaram, S. B., Kumar, R. R., Ramasubramanian, C.,
53 Sivasubramaniam, S., Bjorkum, A. A., Cutajar, J., Berk, M., Trakht, I., Vrdoljak, A., Meira, E.
54 C. M., Eyre, H. A., Gronli, J., Cardinali, D. P., Maercker, A., van de Put, W., Guzder, J.,
55 Bjorvatn, B., Tol, W. A., Acuna-Castroviejo, D., Meudec, M., Morin, C. M., Partinen, M.,
56 Barbui, C., Jordans, M. J. D., Braakman, M. H., Knaevelsrud, C., Pallesen, S., Sijbrandij, M.,
57 Golombek, D. A., Espie, C. A., Cuijpers, P., Agudelo, H. A. M., van der Velden, K., van der
58 Kolk, B. A., Hobfoll, S. E., Deville, W., Gradisar, M., Riemann, D., Axelsson, J., Benitez-King,
59 G., Macy, R. D., Poberezhets, V., Hoole, S. R. H., Murthy, R. S., Hegemann, T., Heinz, A.,
60 Salvage, J., McFarlane, A. C., Keukens, R., de Silva, H., Oestereich, C., Wilhelm, J., von

- 1
2
3
4 Cranach, M., Hoffmann, K., Klosinski, M., Bhugra, D. & Seeman, M. V. (2022). Scientists
5 Against War: A Plea to World Leaders for Better Governance. *Sleep and Vigilance*, 1-6.
6
7 Papola, D., Purgato, M., Gastaldon, C., Bovo, C., van Ommeren, M., Barbui, C. & Tol, W. A. (2020).
8 Psychological and social interventions for the prevention of mental disorders in people
9 living in low- and middle-income countries affected by humanitarian crises. *Cochrane*
10 *Database of Systematic Reviews* 9, CD012417.
11
12 Patel, A. R., Newman, E. & Richardson, J. (2022). A pilot study adapting and validating the
13 Harvard Trauma Questionnaire (HTQ) and PTSD checklist-5 (PCL-5) with Indian women
14 from slums reporting gender-based violence. *BMC Womens Health* 22, 22.
15
16 Plumpton, C. O., Morris, T., Hughes, D. A. & White, I. R. (2016). Multiple imputation of multiple
17 multi-item scales when a full imputation model is infeasible. *BMC Research Notes* 9, 45.
18
19 Priebe, S., Giacco, D. & El-Nagib, R. (2016). *Public Health Aspects of Mental Health Among*
20 *Migrants and Refugees: A Review of the Evidence on Mental Health Care for Refugees,*
21 *Asylum Seekers and Irregular Migrants in the WHO European Region.* WHO Regional
22 Office for Europe: Copenhagen.
23
24 Purgato, M., Carswell, K., Acarturk, C., Au, T., Akbai, S., Anttila, M., Baumgartner, J., Bailey, D.,
25 Biondi, M., Bird, M., Churchill, R., Eskici, S., Hansen, L. J., Heron, P., Ilkkursun, Z., Kilian, R.,
26 Koesters, M., Lantta, T., Nose, M., Ostuzzi, G., Papola, D., Popa, M., Sijbrandij, M.,
27 Tarsitani, L., Tedeschi, F., Turrini, G., Uygun, E., Valimaki, M. A., Wancata, J., White, R.,
28 Zanini, E., Cuijpers, P., Barbui, C. & Van Ommeren, M. (2019). Effectiveness and cost-
29 effectiveness of Self-Help Plus (SH+) for preventing mental disorders in refugees and
30 asylum seekers in Europe and Turkey: study protocols for two randomised controlled
31 trials. *BMJ Open* 9, e030259.
32
33 Purgato, M., Carswell, K., Tedeschi, F., Acarturk, C., Anttila, M., Au, T., Bajbouj, M., Baumgartner,
34 J., Biondi, M., Churchill, R., Cuijpers, P., Koesters, M., Gastaldon, C., Ilkkursun, Z., Lantta,
35 T., Nose, M., Ostuzzi, G., Papola, D., Popa, M., Roselli, V., Sijbrandij, M., Tarsitani, L.,
36 Turrini, G., Valimaki, M., Walker, L., Wancata, J., Zanini, E., White, R., van Ommeren, M. &
37 Barbui, C. (2021). Effectiveness of Self-Help Plus in Preventing Mental Disorders in
38 Refugees and Asylum Seekers in Western Europe: A Multinational Randomized Controlled
39 Trial. *Psychotherapy and Psychosomatics* 90, 403-414.
40
41 Royston, P. (2005). Multiple imputation of missing values: update. *Stata Journal* 5.
42
43 Rytwinski, N. K., Scur, M. D., Feeny, N. C. & Youngstrom, E. A. (2013). The co-occurrence of major
44 depressive disorder among individuals with posttraumatic stress disorder: a meta-
45 analysis. *Journal of Traumatic Stress* 26, 299-309.
46
47 Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T.,
48 Baker, R. & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview
49 (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview
50 for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* 59 Suppl 20, 22-33;quiz 34-57.
51
52 Statacorp (2017). Statacorp.
53
54
55
56
57
58
59
60

- 1
2
3
4 Sterne, J. A., White, I. R., Carlin, J. B., Spratt, M., Royston, P., Kenward, M. G., Wood, A. M. &
5 Carpenter, J. R. (2009). Multiple imputation for missing data in epidemiological and clinical
6 research: potential and pitfalls. *BMJ* 338, b2393.
7
8
9 Stevens, J. (2002). *Applied multivariate statistics for the social sciences*. Lawrence Erlbaum
10 Associates: Mahwah, N.J.
11
12 Tabachnick, B. F., LS (2007). *Using multivariate statistics*. Pearson/Allyn & Bacon: Boston.
13
14 Turrini, G., Tedeschi, F., Cuijpers, P., Del Giovane, C., Kip, A., Morina, N., Nose, M., Ostuzzi, G.,
15 Purgato, M., Ricciardi, C., Sijbrandij, M., Tol, W. & Barbui, C. (2021). A network meta-
16 analysis of psychosocial interventions for refugees and asylum seekers with PTSD. *BMJ*
17 *Global Health* 6.
18
19 UNHCR (2022). Refugee data finder. United Nations High Commissioner for Refugees.
20
21 Uphoff, E., Robertson, L., Cabieses, B., Villalon, F. J., Purgato, M., Churchill, R. & Barbui, C. (2020).
22 An overview of systematic reviews on mental health promotion, prevention, and
23 treatment of common mental disorders for refugees, asylum seekers, and internally
24 displaced persons. *Cochrane Database of Systematic Reviews* 9, CD013458.
25
26
27 White, I. R., Royston, P. & Wood, A. M. (2011). Multiple imputation using chained equations:
28 Issues and guidance for practice. *Statistics in Medicine* 30, 377-99.
29
30
31 White, R. G. & Van der Boor, C. (2021). Enhancing the capabilities of forcibly displaced people: a
32 human development approach to conflict- and displacement-related stressors.
33 *Epidemiology and Psychiatric Sciences* 30, e34.
34
35 World Health Organization, W. H. (2017). *Scalable psychological interventions for people in*
36 *communities affected by adversity*. World Health Organization: Geneva.
37
38 World Health Organization, W. H. (2021a). Self Help Plus (SH+): a group-based stress
39 management course for adults. Generic field-trial version 1.0. World Health Organization:
40 Geneva.
41
42 World Health Organization, W. H. (2021b). *Doing What Matters in Times of Stress*. World Health
43 Organization: Geneva.
44
45 Winkler, J.G., Brandl, E.J., Bretz, H.J., Heinz, A. & Schouler-Ocak, M. (2019) The Influence of
46 Residence Status on Psychiatric Symptom Load of Asylum Seekers in Germany.
47 *Psychiatrische Praxis* 46, 191-199
48
49
50 Wu, Y., Levis, B., Sun, Y., Krishnan, A., He, C., Riehm, K. E., Rice, D. B., Azar, M., Yan, X. W.,
51 Neupane, D., Bhandari, P. M., Imran, M., Chiovitti, M. J., Saadat, N., Boruff, J. T., Cuijpers,
52 P., Gilbody, S., McMillan, D., Ioannidis, J. P. A., Kloda, L. A., Patten, S. B., Shrier, I.,
53 Ziegelstein, R. C., Henry, M., Ismail, Z., Loiselle, C. G., Mitchell, N. D., Tonelli, M., Al-Adawi,
54 S., Beraldi, A., Braeken, A., Buel-Drabe, N., Bunevicius, A., Carter, G., Chen, C. K., Cheung,
55 G., Clover, K., Conroy, R. M., Cukor, D., da Rocha, E. S. C. E., Dabscheck, E., Daray, F. M.,
56 Douven, E., Downing, M. G., Feinstein, A., Ferentinos, P. P., Fischer, F. H., Flint, A. J.,
57 Fujimori, M., Gallagher, P., Gandy, M., Goebel, S., Grassi, L., Harter, M., Jenewein, J., Jette,
58 N., Juliao, M., Kim, J. M., Kim, S. W., Kjaergaard, M., Kohler, S., Loosman, W. L., Lowe, B.,
59
60

1
2
3
4 Martin-Santos, R., Massardo, L., Matsuoka, Y., Mehnert, A., Michopoulos, I., Misery, L.,
5 Navines, R., O'Donnell, M. L., Ozturk, A., Peceliuniene, J., Pintor, L., Ponsford, J. L., Quinn,
6 T. J., Reme, S. E., Reuter, K., Rooney, A. G., Sanchez-Gonzalez, R., Schwarzbald, M. L.,
7 Senturk Cankorur, V., Shaaban, J., Sharpe, L., Sharpe, M., Simard, S., Singer, S., Stafford,
8 L., Stone, J., Sultan, S., Teixeira, A. L., Tiringier, I., Turner, A., Walker, J., Walterfang, M.,
9 Wang, L. J., White, J., Wong, D. K., Benedetti, A. & Thombs, B. D. (2020). Probability of
10 major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic
11 interviews controlling for Hospital Anxiety and Depression Scale - Depression subscale
12 scores: An individual participant data meta-analysis of 73 primary studies. *Journal of*
13 *Psychosomatics Research* 129, 109892.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1. Participants characteristics at baseline

Means	M (SD)
Age (years)	31.797 (9.521)
Years of education	9.556 (4.424)
Months in host country	40.636 (32.599)
Host country GDP in thousand dollars (2019)	
Austria	58,091.3
Finland	50,321.6
Germany	55 652.9
Italy	44,334.2
Turkey	27,144.2
UK	49,070.3
Number of SH+ sessions	1.434 (2.015)
GHQ	5.737 (2.230)
PHQ-9	7.232 (5.230)
HTQ Total score	5.795 (4.349)
HTQ Lack of basic needs	1.277 (1.210)
HTQ Violence and abuse	1.753 (2.085)
HTQ Being close to death	2.766 (1.947)
Frequencies	n/N (%)
Gender (Female)	538/1,101 (48.86%)
Unemployed	418/1,099 (38.03%)
Country of origin	
Nigeria	114/1,100 (10.36%)
Syria	758/1,100 (68.91%)
Iraq	94/1,100 (8.55%)
Other	134/1,100 (12.18%)
Travel duration	
Less than one month	277 (40.44%)
One to three months	203 (29.64%)
More than three months	205 (29.93%)
Study	
Western Europe	459 (41.69%)
Turkey	642 (58.31%)

M: mean; SD: standard deviation; SH+: Self-Help Plus; GDP: Gross Domestic Product; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ: Harvard Trauma Questionnaire

Table 2. Determinants of mental disorder development: logistic regression analyses

	Unadjusted regressions		Adjusted regressions			
			With HTQ total score		With HTQ factors	
	Odds Ratio (95% CI)	P-value	Odds Ratio (95% CI)	P-value	Odds Ratio (95% CI)	P-value
Age in years	1.001 (0.986; 1.015)	0.920	0.996 (0.980; 1.013)	0.672	0.998 (0.981; 1.015)	0.830
Female gender	1.318 (0.987; 1.759)	0.061	0.962 (0.680; 1.360)	0.825	0.977 (0.689; 1.385)	0.896
Years of education	0.972 (0.941; 1.004)	0.088	0.959 (0.923; 0.997)	0.034	0.959 (0.922; 0.996)	0.032
Unemployed	0.938 (0.702; 1.253)	0.664	1.171 (0.832; 1.647)	0.365	1.180 (0.836; 1.664)	0.347
Country of origin:						
Syria	Reference category	<0.001	Reference category	0.017	Reference category	0.006
Iraq	1.180 (0.710; 1.960)		1.592 (0.807; 3.142)		1.672 (0.840; 3.325)	
Nigeria	0.209 (0.101; 0.432)		0.376 (0.142; 0.997)		0.318 (0.118; 0.861)	
Other	0.403 (0.240; 0.674)		0.577 (0.273; 1.222)		0.519 (0.241; 1.114)	
Travel duration (months)						
Less than one month	Reference category	<0.001	Reference category	0.057	Reference category	0.057
One to three months	0.616 (0.398; 0.952)		0.653 (0.414; 1.028)		0.658 (0.417; 1.041)	
More than three months	0.312 (0.198; 0.494)		0.512 (0.267; 0.983)		0.502 (0.260; 0.969)	
Months in host country	1.002 (0.997; 1.007)	0.503	1.000 (0.994; 1.006)	0.943	1.000 (0.994; 1.006)	0.973
Turkish trial	2.058 (1.513; 2.800)	<0.001	1.795 (0.997; 3.235)	0.051	1.802 (0.998; 3.253)	0.051
Number of SH+ sessions	0.888 (0.827; 0.953)	0.001	0.853 (0.788; 0.923)	<0.001	0.850 (0.785; 0.921)	<0.001

GHQ-12	1.188 (1.114; 1.268)	<0.001	1.117 (1.036; 1.204)	0.004	1.119 (1.038; 1.206)	0.003
PHQ-9	1.071 (1.042; 1.100)	<0.001	1.061 (1.026; 1.097)	0.001	1.061 (1.026; 1.097)	0.001
HTQ Total score	1.002 (0.970; 1.036)	0.883	1.051 (1.008; 1.096)	0.020	-	-
HTQ Factor 1 - Lack of basic needs	0.999 (0.890; 1.122)	0.988	-	-	1.007 (0.863; 1.176)	0.928
HTQ Factor 2 - Violence and abuse	0.995 (0.929; 1.066)	0.894	-	-	1.163 (1.041; 1.299)	0.008
HTQ Factor 3 - Being close to death	1.018 (0.947; 1.095)	0.769	-	-	0.978 (0.888; 1.077)	0.650

CI: Confidence Interval; SH+: Self-Help Plus; GDP: Gross Domestic Product; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ: Harvard Trauma Questionnaire

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey: participant-level analysis of two large prevention studies

Corrado Barbui¹, Federico Tedeschi¹, Ceren Acarturk², Minna Anttila³, Teresa Au⁴, Josef Baumgartner⁵, Ken Carswell⁴, Rachel Churchill⁶, Pim Cuijpers^{7,8}, Eirini Karyotaki⁷, Thomas Klein⁹, Markus Koesters⁹, Tella Lantta³, Michela Nosè¹, Giovanni Ostuzzi¹, Massimo Pasquini¹⁰, Eleonora Prina¹, Marit Sijbrandij⁷, Lorenzo Tarsitani¹⁰, Giulia Turrini¹, Ersin Uygun¹¹, Maritta Välimäki^{3,12}, Lauren Walker¹³, Johannes Wancata⁵, Ross G. White¹⁴, Marianna Purgato¹

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

¹ WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, Department of Neuroscience, Biomedicine and Movement Sciences, Section of Psychiatry, University of Verona, Verona 37134, Italy

² Department of Psychology, Koc University, Istanbul, Turkey

³ University of Turku, Department of Nursing Science, 20014 University of Turku, Finland

⁴ Department of Mental Health and Substance Use, World Health Organization, Geneva, Switzerland

⁵ Clinical Division of Social Psychiatry, Department of Psychiatry and Psychotherapy, Medical University of Vienna, Vienna, Austria

⁶ Centre for Reviews and Dissemination, University of York, York, UK

⁷ Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health research institute, Vrije Universiteit Amsterdam, WHO Collaborating Centre for Research and Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam

⁸ Babeş-Bolyai University, International Institute for Psychotherapy, Cluj-Napoca, Romania

⁹ Department of Psychiatry II, Ulm University, Ulm, Germany

¹⁰ Department of Human Neurosciences, Sapienza University of Rome, Italy

¹¹ Trauma and Disaster Mental Health, Istanbul Bilgi University, Eyüpsultan/Istanbul, Turkey

¹² Xiangya Nursing School, Central South University, Xiangya Center for Evidence-Based Practice and Healthcare Innovation, 172 Tongzipo Road, Changsha, Hunan 410013, China

¹³ Department of Health Sciences, University of York, York, UK

¹⁴ School of Psychology, Queen's University Belfast, Belfast, UK

Correspondence to:

Professor Corrado Barbui, WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, Ospedale Policlinico GB Rossi, Piazzale Scuro 10, 37100 Verona, Italy. Email: corrado.barbui@univr.it

Word count: 4,472

For Peer Review

Abstract

Background: In asylum seekers and refugees, the frequency of mental disorders, such as depression, anxiety, and post-traumatic stress disorder, is higher than the general population, but there is a lack of data on risk factors for the development of mental disorders in this population.

Aim: This study investigated the risk factors for mental disorder development in a large group of asylum seekers and refugees resettled in high- and middle-income settings.

Methods: Participant-level data from two randomized prevention studies involving asylum seekers and refugees resettled in Western European countries and in Turkey were pooled. The two studies randomized participants with psychological distress, but without a diagnosis of mental disorder, to the Self-Help Plus psychological intervention or enhanced care as usual. At baseline, exposure to potentially traumatic events was measured using the Harvard Trauma Questionnaire-part I, while psychological distress and depressive symptoms were assessed with the General Health Questionnaire and the Patient Health Questionnaire. After three and six months of follow-up, the proportion of participants who developed a mental disorder was calculated using the Mini International Neuropsychiatric Interview.

Results: A total of 1,101 participants were included in the analysis. At three- and six-month follow-up the observed frequency of mental disorders was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. After controlling for confounders, logistic regression analysis showed that participants with a lower education level ($p = 0.034$), a shorter duration of journey ($p = 0.057$), and arriving from countries with war-related contexts ($p = 0.017$), were more at risk of developing mental disorders. Psychological distress ($p = 0.004$), depression ($p = 0.001$), and

1
2
3
4 exposure to potentially traumatic events ($p = 0.020$) were predictors of mental disorder
5
6 development.
7
8

9 **Conclusions:** This study identified several risk factors for the development of mental disorders
10
11 in asylum seekers and refugees, some of which may be the target of risk reduction policies. The
12
13 identification of asylum seekers and refugees at increased risk of mental disorders should guide
14
15 the implementation of focused preventative psychological interventions.
16
17
18
19

20
21
22 **Key-words:** asylum seekers, refugees, migrants, mental health, mental disorders, high-income
23
24 countries, low-income countries
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

According to the last United Nations Refugee Agency data, despite COVID-related movement restrictions, at the end of 2020 more than 82 million people were forcibly displaced as a result of persecution, conflict, violence, and other human rights violations (UNHCR, 2022). Of these, more than 30 million were asylum seekers or refugees. While Turkey continued to host the largest number of refugees, most of whom were Syrian refugees, 10% of all the world's refugees were resettled in Europe at the end of 2020 (UNHCR, 2022). These numbers are expected to further increase, as recent estimates have suggested that over 6.5 million refugees from Ukraine have crossed to Poland, Hungary, Romania, Moldova and other countries (UNHCR, 2022).

Despite heterogeneity between the studies and the populations of refugees and asylum seekers assessed (Giacco, 2019), a large body of evidence consistently showed that the frequency of mental disorders in refugees and asylum seekers is increased as compared with the general population. Existing World Health Organization (WHO) estimates suggest a prevalence of 13.0% for mild forms of depression, anxiety, and post-traumatic stress disorder (PTSD), and 4.0% for moderate forms (Charlson *et al.*, 2019). Other reviews of prevalence studies found adult refugee and asylum seekers have high and persistent rates of depression and PTSD, while the prevalence of anxiety disorders and psychosis are more comparable to findings from general populations (Blackmore *et al.*, 2020, Henkelmann *et al.*, 2020).

The increased frequency of mental disorders in refugees and asylum seekers may be related to the refugee experience, which is characterized by loss of homes, hopes, possessions, and disruption of personal, family, and professional life projects. In addition, a wide range of physical, psychological, and psychosocial problems associated with adversities may occur, such as bombings, threats, captivity, torture, injury, witnessing death or injury of loved ones, discrimination, economic stress, and uncertainty about the future (Priebe *et al.*, 2016).

1
2
3
4 Interestingly, while these factors have been studied as determinants of poor mental health
5 outcomes, or as determinants of psychological distress (Jannesari *et al.*, 2020), to our knowledge
6 they have never been investigated in this population as risk factors for the development of
7 mental disorders considered as full-blown diagnostic entities (Gleeson *et al.*, 2020, Hajak *et al.*,
8 2021). Knowledge of determinants of the development of mental disorders is particularly
9 important to implement policies aiming at decreasing exposure to such determinants, to
10 improve identification of refugees and asylum seekers at increased risk of developing mental
11 disorders, and to plan the provision of psychological interventions to individuals and
12 communities at risk.

13
14
15
16
17
18
19
20
21
22
23
24
25 Aiming to test the effect of Self Help Plus (SH+), a low intensity, group-based, self-help
26 psychological intervention recently developed by the WHO (Epping-Jordan *et al.*, 2016, World
27 Health Organization, 2021a), in reducing the frequency of mental disorders, we conducted two
28 large randomized prevention trials in asylum seekers and refugees, one in western Europe and
29 another in Turkey (Purgato *et al.*, 2019). The studies followed the same research protocol and
30 were conducted in parallel (Acarturk *et al.*, 2022, Purgato *et al.*, 2021). Both the Western
31 European and Turkey studies showed evidence of an effect of SH+ in preventing the onset of
32 mental disorders, but differences were observed between the studies. The effect was much
33 more pronounced for the Turkey study where efficacy (i.e. reducing the frequency of any mental
34 disorder) was observed at 6 months, compared to the Western European study where a
35 preventative effect was only found post-intervention and not after six months (Acarturk *et al.*,
36 2022, Purgato *et al.*, 2021). As these are the only two studies that enrolled asylum seekers and
37 refugees without any mental disorders at baseline, and that assessed the frequency of mental
38 disorders at follow-up as primary outcome (Papola *et al.*, 2020), they offered a unique
39 opportunity to prospectively investigate the risk factors for the onset of mental disorders in a
40 large group of asylum seekers and refugees.

Methods

Participants and measures

Participant-level data from two randomized prevention trials involving asylum seekers and refugees resettled in Western European countries (Austria, Finland, Germany, Italy, and UK) and in Turkey were pooled (Acarturk *et al.*, 2022, Purgato *et al.*, 2021). In both studies, participants were randomly assigned to the SH+ psychological intervention, consisting of SH+ combined with enhanced care as usual (ECAU), or to ECAU only. The two studies were conducted in parallel following the same study design (Purgato *et al.*, 2019). The protocol of the present study was registered within the Open Science Framework (<https://osf.io/37h5n>).

In both studies, participants were included if they met the following criteria: a) aged 18 years or older; b) able to speak and understand Arabic, Dari, English, or Urdu; c) being under temporary protection with a refugee or asylum seeker status; d) experiencing psychological distress, as shown by a score of 3 or more on the 12-item dichotomously-scored General Health Questionnaire (el-Rufaie and Daradkeh, 1996, Goldberg *et al.*, 1998, Kilic *et al.*, 1997); e) having completed oral and written informed consent to enter the study. Exclusion criteria were: a) presence of any mental disorder according to the Mini International Neuropsychiatric Interview (M.I.N.I.), a brief structured diagnostic interview for the major psychiatric disorders in DSM-III-R, DSM-IV and DSM-5 and ICD-10. (Kadri *et al.*, 2005, Sheehan *et al.*, 1998); b) evidence of acute medical conditions contraindicating study participation; c) evidence of imminent suicide risk, or suicide risk scored as “moderate or high” on the M.I.N.I.; d) signs of impaired decision-making capacity emerging from responses during the clinical interview.

The M.I.N.I. was administered before random allocation, in order to exclude participants with a mental disorder, and at three and six months of follow-up, in order to calculate the proportion

1
2
3
4 of participants who developed a mental disorder. Exposure to potentially traumatic events was
5
6 measured at baseline using the Harvard Trauma Questionnaire-part I (HTQ) (Mollica *et al.*,
7
8 1992). The HTQ-part I covers a variety of trauma events that may affect refugee mental health,
9
10 and the scoring represents the number of different types of traumatic events experienced by
11
12 the participants (higher score is associated with high number of traumatic events). In addition
13
14 to using the HTQ total score, we identified subtypes of traumatic events performing a Principal
15
16 Components analysis (PCA).
17
18
19

20
21 Mental health symptoms were measured using the following validated instruments. The GHQ-
22
23 12 questionnaire was used to measure psychological distress (el-Rufaie and Daradkeh, 1996,
24
25 Goldberg *et al.*, 1998, Kilic *et al.*, 1997), while PTSD was assessed with the PTSD Checklist for
26
27 DSM-5 (PCL-5), a 20-item questionnaire that measures overall PTSD symptoms (score zero to
28
29 80) and symptoms by cluster (intrusions, avoidance, negative changes in thoughts and mood,
30
31 and changes in arousal), with higher scores indicating higher levels of PTSD symptoms
32
33 (Blanchard *et al.*, 1996). Levels of self-reported depression symptoms were measured with the
34
35 Patient Health Questionnaire, nine-item version (PHQ-9) which has a four-point scale (score 0
36
37 to 27) (Kroenke *et al.*, 2001). All measures were collected at baseline before random allocation,
38
39 and after three and six months of follow-up. Assessors were trained in the administration of
40
41 rating scales, instructed on how to perform follow-up assessments while preserving effective
42
43 masking, and assisted by cultural mediators when needed.
44
45
46
47

48 *Interventions*

49
50
51 The SH+ intervention was developed by the WHO, as described elsewhere (Acarturk *et al.*, 2022,
52
53 Purgato *et al.*, 2021), and is now publicly available (World Health Organization, 2021a). SH+
54
55 consists of a pre-recorded audio course complemented with an illustrated self-help book. The
56
57 book has been recently updated and published by WHO as *Doing What Matters in Times of*
58
59
60

1
2
3
4 Stress (World Health Organization, 2021b). All SH+ materials were adapted for the cultural
5
6 groups included in the studies. The SH+ pre-recorded audio material was delivered across five
7
8 2-hour sessions to groups of up to 30 people. The audio material imparts key information about
9
10 stress management and guides participants through individual exercises and small group
11
12 discussions. To augment the audio recordings, an illustrated self-help book reviews all essential
13
14 content and concepts. The SH+ intervention was fully delivered in the language of participants
15
16 by trained facilitators with a migration background, who were native speakers of the target
17
18 languages. ECAU, provided both to the experimental and control group, consisted of routinely
19
20 delivered social support and/or care according to local regulations. Participants in the ECAU arm
21
22 received the same baseline and follow-up assessments of the intervention arm, according to the
23
24 study schedule (around three and six months after randomization), information about freely
25
26 available health and social services, and links to community networks providing support to
27
28 refugees and asylum seekers.
29
30
31
32

33 34 *Statistical analysis*

35
36 Descriptive statistics (mean and SD for continuous variables and absolute numbers and
37
38 percentages for dichotomous variables) were computed on sociodemographic, premigration,
39
40 migration and postmigration variables at baseline, and for clinical variables.
41
42
43

44 Participants who met criteria for any mental disorder on the M.I.N.I. at three or six-month
45
46 follow-up were considered cases with a mental disorder. Multiple imputation was adopted to
47
48 address the issue of missing data in all the variables included in the model. In particular, in case
49
50 of missing data at the M.I.N.I. in one timepoint, imputation was performed on single M.I.N.I.
51
52 values at three and six months. In case of missing values on continuous clinical measures, the
53
54 imputation was performed on single item scales. Specifically, imputation followed the approach
55
56 reported by Plumpton and colleagues (Plumpton *et al.*, 2016), that is we used scale totals within
57
58
59
60

1
2
3
4 prediction equations and, for imputations of responses to individual scale items, we additionally
5 included the responses to the other scale items, using the “ice” Stata routine (Royston, 2005,
6 Sterne *et al.*, 2009), and considering single-item scores as ordered categorical variables. M.I.N.I.
7 values at the two time-points were used in the prediction equations of regression predictors,
8 and upper and lower bounds were set for continuous variables with missing values as
9 appropriate. The number of imputed samples was determined by following the rule of thumb
10 suggested by White and colleagues, i.e.: “at least equal to the percentage of incomplete cases”
11 (White *et al.*, 2011). We rounded such number to the nearest multiple of 10 above.

12
13
14
15
16
17
18
19
20
21
22
23 In order to identify subtypes of traumatic events, we performed a PCA on tetrachoric
24 correlations of HTQ items, with Quartimin oblique rotation to allow for between-factor non-null
25 correlations (Tabachnick, 2007), and summing items with loadings above 0.40 on the same
26 factor to create scores for the regression model (Stevens, 2002). The number of factors was
27 determined by adopting Kaiser's rule (Kaiser, 1960), i.e. using the cut-off scores of 1 for the
28 eigenvalues.
29
30
31
32
33
34
35
36

37 In order to investigate predictors of the development of mental disorders, unadjusted and
38 adjusted logistic regression analyses were performed, using the frequency of participants with
39 a mental disorder in at least one timepoint as binary dependent variable. The following
40 independent variables were inserted into the model: age (years), gender (men, women),
41 education (years), unemployment (yes/no), country of origin (Syria, Iraq, Nigeria, other
42 countries), length of journey (below one month, between one and three months, above three
43 months), study (Western Europe versus Turkey), length of stay in the resettlement country
44 (months), HTQ (total score at baseline), GHQ-12 (total score at baseline), PCL-5 (total score at
45 baseline), PHQ-9 (total score at baseline), number of SH+ sessions received (zero to five).
46
47
48
49
50
51
52
53
54
55
56
57 Considering that PTSD and depression are known to be interrelated experiences following
58
59
60

1
2
3
4 trauma (Rytwinski *et al.*, 2013), and considering that the PCL-5 largely overlaps with the HTQ
5
6 (Patel *et al.*, 2022), and that a substantial overlap exists between the PCL-5 and the PHQ-9
7
8 (Dabrowski CL, 2020), we excluded the PCL-5 from the final model, but it was included (total
9
10 score at baseline) in the imputation model.
11
12

13
14 In order to check the robustness of the analysis, we re-ran the model after excluding the arm
15
16 receiving SH+. As a further sensitivity analysis, in order to estimate within-centre effects, we
17
18 performed a model including a fixed effect for recruiting centre. As a subgroup analysis, we re-
19
20 ran the logistic regression analysis separately for each study sample (western Europe versus
21
22 Turkey). All analyses were performed using Stata 17 (Statacorp, 2017).
23
24
25
26
27

28 **Results**

29 *Participants*

30
31
32
33
34 The two prevention trials randomized a total of 1,101 participants. After six months, 249
35
36 participants (22.6%) were not available for follow-up assessments, for the reasons reported in
37
38 Figure 1. Participant characteristics at baseline are presented in Table 1. Almost half were
39
40 female, with a mean age of 32 (SD 9.521) years, and a mean education of 10 years (SD 4.424).
41
42 Slightly less than 40% was unemployed. The migration journey lasted more than three months
43
44 in around one third of participants, in another one third it lasted between one and three months,
45
46 while in the remaining 40% the host country was reached in less than one month. On average,
47
48 the mean length of stay in the resettlement country was slightly more than three years. Most of
49
50 the participants were from Syria, Nigeria and Iraq (Table 1). Participants attended a mean
51
52 number of 1.4 (SD 2.015) SH+ sessions. The mean (SD) baseline scores on the measures of
53
54 interest are reported in Table 1.
55
56
57
58
59
60

Frequency and determinants of mental disorders

At three- and six-month follow-up the observed frequency of mental disorders, as measured with the M.I.N.I., was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. The majority of detected mental disorders were major depressive disorders (9.5% and 20.3% after three and six months), PTSD (3.2% and 6.7%), and anxiety disorders (2.1% and 4.1%) (Figure 1).

The PCA model found a 3-factor solution where the set of items was exhaustive and mutually exclusive, with the HTQ items grouped as follows: factor 1 - lack of basic needs: lack of food or water, ill health without access to medical care, lack of shelter; factor 2 - violence and abuse: imprisonment, serious injury, brain washing, rape or sexual abuse, forced isolation from others, forced separation from family members, torture, other (e.g. domestic violence;) factor 3 - being close to death: combat situation, being close to death, murder of family or friend, unnatural death of family or friend, murder of stranger(s), lost or kidnapped. The factor loadings of the PCA-model with Quartimin oblique rotation are reported in the Supplemental Materials.

The results of unadjusted and adjusted logistic regression analyses investigating factors associated with the development of mental disorders are presented in Table 2. Participants with lower education level ($p = 0.034$), a shorter duration of journey ($p = 0.057$), and arriving from countries with war-related contexts (Iraq, Syria) ($p = 0.017$), were those more at risk of developing mental disorders. In terms of mental health symptoms and exposure to traumatic events, psychological distress ($p = 0.004$), depressive symptoms ($p = 0.001$), and HTQ total score ($p = 0.020$) at baseline were predictors of mental disorder development, as was the HTQ factor violence and abuse (imprisonment, serious injury, brain washing, rape or sexual abuse, forced isolation from others, forced separation from family members, torture, domestic violence) ($p =$

1
2
3
4 0.008). In terms of post-migration factors, the number of SH+ sessions ($p < 0.001$) was inversely
5 associated with the risk of developing mental disorders (Table 2).
6
7

8
9 Secondary logistic regression analyses including only the ECAU arms of the two studies, and
10 analysing the two studies separately, confirmed the role of mental health symptoms and
11 exposure to traumatic events as risk factors for mental disorders (Supplemental Materials).
12
13 However, due to lower statistical power, some factors lost significance despite a similar or even
14 higher estimated effect, as for example the HTQ in the model restricted to the western European
15 study, or years of education and travel duration in the model restricted to the ECAU sample, and
16 GHQ-12 in the model restricted to the study conducted in Turkey (Supplemental Materials).
17
18
19
20
21
22
23
24
25

26 Consistently with the results of the two studies, the number of SH+ sessions was inversely
27 associated with the frequency of mental disorders only in the study conducted in Turkey
28 Supplemental Materials). The inclusion of recruiting centres as predictor variables revealed that,
29 in comparison with participants recruited in Istanbul, participants recruited in Vienna were less
30 likely to develop a mental disorder (Supplemental Materials).
31
32
33
34
35
36
37
38
39
40

41 **Discussion**

42
43 To the best of our knowledge, this is the first prospective study that examined risk factors for
44 the onset of mental disorders among asylum seekers and refugees without a mental disorder at
45 baseline. We showed the significant influence of socio-demographic, clinical, and contextual
46 factors, including potentially traumatic events, on the development of mental disorders in
47 asylum seekers and refugees resettled in Western European countries and in Turkey.
48
49
50
51
52
53
54

55 Overall, one in four refugees and asylum seekers developed a diagnosable mental disorder over
56 a period of six months. Interestingly, the most frequently reported diagnostic group was
57
58
59
60

1
2
3
4 depression, followed by PTSD and anxiety disorders. These figures, derived from two
5
6 intervention studies, cannot be compared with those from epidemiological studies, which
7
8 generally show similar frequencies for depression and PTSD, or slightly higher frequencies for
9
10 depression, depending on the population surveyed and the study setting (Blackmore *et al.*, 2020,
11
12 Charlson *et al.*, 2019, Henkelmann *et al.*, 2020, Hoell *et al.*, 2021). In migrants exposed to armed
13
14 conflict, by contrast, the frequency of PTSD was found to be higher than depression (31% vs
15
16 25%) (Mesa-Vieira *et al.*, 2022).
17
18
19

20
21 The finding that educational level is inversely associated with the development of mental
22
23 disorders expands previous data collected in the general population showing that higher
24
25 educational level seems to have a protective effect against anxiety and depression symptoms,
26
27 and against common mental disorders in general (Araya *et al.*, 2003, Bjelland *et al.*, 2008). In the
28
29 general population, poor education has been suggested to be a marker of lack of opportunities
30
31 and resources, including material and psychological resources, and a marker of childhood
32
33 adversity (Araya *et al.*, 2003). In asylum seekers and refugees, these factors may directly
34
35 contribute to the emergence of mental disorders, or may interfere with the coping skills required
36
37 to deal with all the adversities associated with the migration and resettlement process, which,
38
39 in turn, may increase the risk of developing mental disorders (Kirmayer *et al.*, 2011).
40
41
42

43
44 Consistently with an extensive literature showing that exposure to potentially traumatic events
45
46 represents a risk factor for poor mental health and well-being (Gleeson *et al.*, 2020, Hajak *et al.*,
47
48 2021, Priebe *et al.*, 2016), the present study adds that exposure to potentially traumatic events
49
50 is also a risk factor for the development of mental disorders considered as full-blown diagnostic
51
52 entities. The finding that being displaced from countries with war-related contexts emerged as
53
54 an additional risk factor for mental disorders further corroborates this association. War-related
55
56 contexts imply exposure to multiple and serious traumatic events, and the exposure to combat
57
58
59
60

1
2
3
4 situations creates the risk of witnessing violence and/or death and experiencing physical and
5
6 psychological violence (Crepet *et al.*, 2017). Extensive literature shows that these situations are
7
8 associated with poor mental health outcomes, including PTSD, depression, anxiety, and
9
10 somatization symptoms (Jongedijk *et al.*, 2020, Knipscheer *et al.*, 2015, Nickerson *et al.*, 2021).

11
12
13
14 The present study has limitations and strengths. A first limitation is that the population enrolled
15
16 in the two trials cannot be considered representative of the general population of asylum
17
18 seekers and refugees, as we selected participants scoring above a threshold of psychological
19
20 distress, and we excluded those at suicide risk. During the studies, in addition, around half of
21
22 the participants was exposed to a psychological intervention aimed at preventing the
23
24 development of mental disorders. For these reasons, the overall frequency estimate of mental
25
26 disorders cannot apply to the general population of asylum seekers and refugees resettled in
27
28 Western European countries or in Turkey. A second issue is that we did not calculate a true
29
30 incidence of mental disorders, but only frequency figures at two time points with a structured
31
32 diagnostic interview that might overestimate the true frequency of some mental disorders, such
33
34 as depression (Wu *et al.*, 2020). Third, the SARS-CoV2 pandemic impacted the study procedures,
35
36 because in all the recruiting sites follow-up assessments were conducted using online tools
37
38 instead of face-to-face meetings. Although assessors were trained in the administration of rating
39
40 scales, instructed on how to perform online follow-up assessments, and assisted by cultural
41
42 mediators when needed, it is unknown whether this may have impacted the responses of
43
44 participants to the instruments. Related to this, even though several studies documented that a
45
46 careful and culturally appropriate use of available instruments is feasible and allows a
47
48 standardization of the screening process and a systematic recognition of psychological distress
49
50 and psychiatric diagnoses (Acarturk *et al.*, 2021) we acknowledge that formal studies on use of
51
52 these tools in refugee groups are lacking. Fourth, despite a growing body of literature showed
53
54 that the duration of the asylum procedure is an important risk factor for mental health
55
56
57
58
59
60

1
2
3
4 conditions (Laban et al. 2004; Winkler et al. 2019), we acknowledge that this information was
5
6 not collected. However, months in host country, which may be considered a proxy of length of
7
8 asylum procedure, was included in the model but did not emerge as a significant factor.
9

10
11
12 Despite these limitations, there are strengths that should be emphasised. The main strength is
13
14 the exclusion of participants with a mental disorder at baseline, and the choice of a dependent
15
16 variable that is fully consistent with a preventative design, namely the frequency of mental
17
18 disorders at follow-up, assessed with validated measures. This design allowed to prospectively
19
20 investigate the determinants of mental disorders rather than of poor mental health, as previous
21
22 studies have done in populations who might already be with mental disorders at baseline (Priebe
23
24 *et al.*, 2016). Another strength is a sample size of over a thousand of asylum seekers and
25
26 refugees, resettled in a variety of western European sites and in Turkey. This aspect is of
27
28 relevance not only in terms of statistical power, but also in terms of generalisability and
29
30 applicability of study findings to different types of reception settings in high-income and middle-
31
32 income countries. Lastly, despite an attrition rate of around 20%, a follow-up assessment of six
33
34 months is noteworthy in such a difficult-to-follow population, who is often moved from one
35
36 reception site to another, and may not perceive mental health as a priority, having a number of
37
38 other challenging concerns such as housing, unstable working conditions, management of visa
39
40 issues, safety of family members, fear of being returned to home country, plans to move to
41
42 another country or to another location.
43
44
45
46
47

48
49 The present study has important practical implications. The finding of a positive relationship
50
51 between exposure to potentially traumatic events and risk of mental disorder development
52
53 suggests a pressing need of developing policies aiming to decrease exposure to such traumatic
54
55 experiences after resettlement. Host countries may have opportunities to decrease exposure to
56
57 situations such as material and economic hardship that could affect integrity, independence,
58
59
60

1
2
3
4 dignity and well-being (financial strain), social hardship due to loss of status (social strain),
5
6 feelings of inadequacy in relation with specific skills needed in the host-country to successfully
7
8 function in daily life (competency strain), experiences of unfair treatment on the basis of
9
10 prejudice (perceived discrimination). This ambitious goal may be achieved by implementing
11
12 reception conditions that optimize internationally recognized minimal quality standards. For
13
14 example, standards for the reception of applicants for international protection have been
15
16 established by Directive 2013/33/EU of the European Parliament. The Directive clearly reports
17
18 that national authorities should ensure that reception modalities are specifically designed to
19
20 meet the needs of persons requiring international protection, including legal assistance,
21
22 document provision, material support, links with local communities, freedom of movement,
23
24 information about labor market access, vocational training, social support. Health care, including
25
26 mental health care, is also mentioned as a key intervention where needed. It would be important
27
28 to ensure that efforts to support forcibly displaced people are coordinated across the different
29
30 layers of the social environments in which they are hosted, i.e. at the level of the individual, their
31
32 family, the community, and the institutions that have governance responsibility for their care
33
34 and support (White and Van der Boor, 2021).
35
36
37
38
39
40

41 In addition to implications in terms of risk reduction policies, the present study has implications
42
43 in terms of provision of psychological interventions aimed at preventing the development of
44
45 mental disorders (Miller *et al.*, 2021). As the implementation of preventative psychological
46
47 interventions to the whole population of asylum seekers and refugees may not be sustainable
48
49 by host countries, national authorities may consider to offer psychological interventions to at-
50
51 risk populations. The finding that persons with lower education level, a shorter duration of
52
53 journey, arriving from countries with war-related contexts (Iraq, Syria), and with high level of
54
55 psychological distress and depressive symptoms, are those more at risk of developing mental
56
57 disorders, appears to identify a target population for focused psychological prevention
58
59
60

1
2
3
4 interventions. The WHO has recently developed a number of low-intensity psychological
5
6 interventions that may be scaled up as public health strategies to address mental disorders and
7
8 psychological distress in refugee populations exposed to ongoing adversities (World Health
9
10 Organization, 2017). In addition to WHO interventions, other psychological treatments have
11
12 been shown to be effective in alleviating psychological symptoms in asylum seekers and
13
14 refugees (Turrini *et al.*, 2021, Uphoff *et al.*, 2020), but a preventative effect has been
15
16 investigated for SH+ only. The present results, by showing an inverse association between
17
18 number of SH+ sessions and risk of developing a mental disorder, further corroborate the value
19
20 of SH+ as a prevention intervention. The finding that trauma exposure stands out as a predictor
21
22 for development of mental disorders would additionally suggest that interventions may need to
23
24 include a focus on traumatic memories and experiences, in addition to being focused on low
25
26 mood.
27
28
29
30

31
32 In parallel with the provision of psychological interventions to at-risk populations, national
33
34 authorities should ensure regular access to such interventions supporting at risk populations to
35
36 engage through reducing barriers that might prevent or limit access or use (Fuhr *et al.*, 2019).
37
38 This can be achieved by optimizing access (e.g. accessible services and supports) to a range of
39
40 interventions depending on need (e.g. stepped care), and by using different delivery
41
42 mechanisms (e.g. digital or peer delivered support).
43
44
45

46
47 As the number of persons in need of protection is likely to substantially increase globally, driven
48
49 by long-lasting wars as well as by new conflicts such as the Russo-Ukrainian war which broke out
50
51 recently (Barbui *et al.*, 2022, Pandi-Perumal *et al.*, 2022), national authorities are urged to
52
53 develop reception and resettlement programs meeting the needs of this vulnerable group.
54
55 These programs should be designed and implemented attempting to decrease the risk of post-
56
57 migration stressors that may contribute to worsening the mental health of a population already
58
59
60

1
2
3
4 exposed to potentially traumatic experiences before and during the migration process.
5
6 Evidence-based focused psychological support should be an important program component to
7
8 be delivered to asylum seekers and refugees at-risk of developing a mental disorder.
9
10
11
12

13 **Contributors** CB, MP, and FT conceived the study. FT statistically analysed the data. CB, GO
14 and FT accessed and verified the data. CB wrote the first draft of the manuscript with input
15 from FT and MP. All authors contributed to the conception and design of the study, data
16 interpretation and manuscript revision. All authors read and approved the submitted
17 manuscript and had final responsibility for the decision to submit for publication. CB was the
18 overall guarantor of the content.
19
20
21

22 **Funding** This work was supported by the European Commission (grant agreement No. 779255;
23 “RE-DEFINE: Refugee Emergency: DEFining and Implementing Novel Evidence-based
24 psychosocial interventions”).
25
26
27

28 **Disclaimer** The authors alone are responsible for the views expressed in this article, and they
29 do not necessarily represent the views, decisions, or policies of the institutions with which they
30 are affiliated.
31
32
33

34 **Competing interests** None declared.
35
36
37

38 **Ethics approval** This was a secondary analysis of deidentified participant data. The initial study
39 was approved by the WHO Research Ethics Review Committee, and from the Ethics
40 Committees of all participating sites. Before participation, a written informed consent was
41 obtained from each participant, allowing investigators to use anonymised data for future
42 analysis. Participants gave informed consent to participate in the study before taking part.
43
44
45

46 **Data availability statement** Data collected in the RE-DEFINE study are stored in the online
47 repository EUDAT B2SHARE (study in Western Europe:
48 <https://b2share.eudat.eu/records/fa7264d624364683830ff37acee01c04>; study in Turkey:
49 <http://doi.org/10.23728/b2share.8ac4f28d2415413e89de7847c05471fc>). The reuse of data
50 will be offered only upon motivated request, which will undergo the scrutiny of the RE-DEFINE
51 General Assembly.
52
53
54
55
56
57
58
59
60

References

- [Acarturk, C., McGrath, M., Roberts, B., Ilkkursun, Z., Cuijpers, P., Sijbrandij, M., Sondorp, E., Ventevogel, P., McKee, M. & Fuhr DC. \(2021\) Prevalence and predictors of common mental disorders among Syrian refugees in Istanbul, Turkey: a cross-sectional study. *Social Psychiatry and Psychiatric Epidemiology* 56, 475-84.](#)
- Acarturk, C., Uygun, E., Ilkkursun, Z., Carswell, K., Tedeschi, F., Batu, M., Eskici, S., Kurt, G., Anttila, M., Au, T., Baumgartner, J., Churchill, R., Cuijpers, P., Becker, T., Koesters, M., Lantta, T., Nose, M., Ostuzzi, G., Popa, M., Purgato, M., Sijbrandij, M., Turrini, G., Valimaki, M., Walker, L., Wancata, J., Zanini, E., White, R. G., van Ommeren, M. & Barbui, C. (2022). Effectiveness of a WHO self-help psychological intervention for preventing mental disorders among Syrian refugees in Turkey: a randomized controlled trial. *World Psychiatry* 21, 88-95.
- Araya, R., Lewis, G., Rojas, G. & Fritsch, R. (2003). Education and income: which is more important for mental health? *Journal of Epidemiology and Community Health* 57, 501-5.
- Barbui, C., Purgato, M., Acarturk, C., Churchill, R., Cuijpers, P., Koesters, M., Sijbrandij, M., Valimaki, M., Wancata, J. & White, R. G. (2022). Preventing the mental health consequences of war in refugee populations. *Epidemiology and Psychiatric Sciences* 31, e24.
- Bjelland, I., Krokstad, S., Mykletun, A., Dahl, A. A., Tell, G. S. & Tambs, K. (2008). Does a higher educational level protect against anxiety and depression? The HUNT study. *Social Science and Medicine* 66, 1334-45.
- Blackmore, R., Boyle, J. A., Fazel, M., Ranasinha, S., Gray, K. M., Fitzgerald, G., Misso, M. & Gibson-Helm, M. (2020). The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis. *PLoS Medicine* 17, e1003337.
- Blanchard, E. B., Jones-Alexander, J., Buckley, T. C. & Forneris, C. A. (1996). Psychometric properties of the PTSD Checklist (PCL). *Behavioural Research and Therapeutics* 34, 669-73.
- Charlson, F., van Ommeren, M., Flaxman, A., Cornett, J., Whiteford, H. & Saxena, S. (2019). New WHO prevalence estimates of mental disorders in conflict settings: a systematic review and meta-analysis. *Lancet* 394, 240-248.
- Crepet, A., Rita, F., Reid, A., Van den Boogaard, W., Deiana, P., Quaranta, G., Barbieri, A., Bongiorno, F. & Di Carlo, S. (2017). Mental health and trauma in asylum seekers landing in Sicily in 2015: a descriptive study of neglected invisible wounds. *Conflict and Health* 11, 1.
- Dabrowski CL, L. J., Sherman MF, Kirkhart MW, Breen D (2020). Probable Ptsd, Symptoms of Depression, and an Exploratory Factor Analysis of Item Responses in Canadian Steelworkers After Workplace Traumas. *Crisis, Stress, and Human Resilience: An International Journal* 2.

- 1
2
3
4 el-Rufaie, O. F. & Daradkeh, T. K. (1996). Validation of the Arabic versions of the thirty- and
5 twelve-item General Health Questionnaires in primary care patients. *British Journal of*
6 *Psychiatry* 169, 662-4.
7
8
9 Epping-Jordan, J. E., Harris, R., Brown, F. L., Carswell, K., Foley, C., Garcia-Moreno, C., Kogan, C.
10 & van Ommeren, M. (2016). Self-Help Plus (SH+): a new WHO stress management
11 package. *World Psychiatry* 15, 295-296.
12
13 Fuhr, D. C., Acarturk, C., McGrath, M., Ilkkursun, Z., Sondorp, E., Sijbrandij, M., Ventevogel, P.,
14 Cuijpers, P., McKee, M. & Roberts, B. (2019). Treatment gap and mental health service
15 use among Syrian refugees in Sultanbeyli, Istanbul: a cross-sectional survey. *Epidemiology*
16 *and Psychiatric Sciences* 29, e70.
17
18 Giacco, D. (2019). Identifying the critical time points for mental health of asylum seekers and
19 refugees in high-income countries. *Epidemiology and Psychiatric Sciences* 29, e61.
20
21 Gleeson, C., Frost, R., Sherwood, L., Shevlin, M., Hyland, P., Halpin, R., Murphy, J. & Silove, D.
22 (2020). Post-migration factors and mental health outcomes in asylum-seeking and
23 refugee populations: a systematic review. *European Journal of Psychotraumatology* 11,
24 1793567.
25
26 Goldberg, D. P., Oldehinkel, T. & Ormel, J. (1998). Why GHQ threshold varies from one place to
27 another. *Psychological Medicine* 28, 915-21.
28
29 Hajak, V. L., Sardana, S., Verdeli, H. & Grimm, S. (2021). A Systematic Review of Factors Affecting
30 Mental Health and Well-Being of Asylum Seekers and Refugees in Germany. *Frontiers in*
31 *Psychiatry* 12, 643704.
32
33 Henkelmann, J. R., de Best, S., Deckers, C., Jensen, K., Shahab, M., Elzinga, B. & Molendijk, M.
34 (2020). Anxiety, depression and post-traumatic stress disorder in refugees resettling in
35 high-income countries: systematic review and meta-analysis. *BJPsych Open* 6, e68.
36
37 Hoell, A., Kourmpeli, E., Salize, H. J., Heinz, A., Padberg, F., Habel, U., Kamp-Becker, I., Hohne, E.,
38 Boge, K. & Bajbouj, M. (2021). Prevalence of depressive symptoms and symptoms of post-
39 traumatic stress disorder among newly arrived refugees and asylum seekers in Germany:
40 systematic review and meta-analysis. *BJPsych Open* 7, e93.
41
42 Jannesari, S., Hatch, S. & Oram, S. (2020). Seeking sanctuary: rethinking asylum and mental
43 health. *Epidemiology and Psychiatric Sciences* 29, e154.
44
45 Jongedijk, R. A., Eising, D. D., van der Aa, N., Kleber, R. J. & Boelen, P. A. (2020). Severity profiles
46 of posttraumatic stress, depression, anxiety, and somatization symptoms in treatment
47 seeking traumatized refugees. *Journal of Affective Disorders* 266, 71-81.
48
49 Kadri, N., Agoub, M., El Gnaoui, S., Alami Kh, M., Hergueta, T. & Moussaoui, D. (2005). Moroccan
50 colloquial Arabic version of the Mini International Neuropsychiatric Interview (MINI):
51 qualitative and quantitative validation. *European Psychiatry* 20, 193-5.
52
53 Kaiser, H. F. (1960). The Application of Electronic Computers to Factor Analysis. *Educational and*
54 *Psychological Measurement* 20.
55
56
57
58
59
60

- 1
2
3
4 Kilic, C., Rezaki, M., Rezaki, B., Kaplan, I., Ozgen, G., Sagduyu, A. & Ozturk, M. O. (1997). General
5 Health Questionnaire (GHQ12 & GHQ28): psychometric properties and factor structure of
6 the scales in a Turkish primary care sample. *Social Psychiatry and Psychiatric Epidemiology*
7 32, 327-31.
8
9
10 Kirmayer, L. J., Narasiah, L., Munoz, M., Rashid, M., Ryder, A. G., Guzder, J., Hassan, G., Rousseau,
11 C., Pottie, K., Canadian Collaboration for, I. & Refugee, H. (2011). Common mental health
12 problems in immigrants and refugees: general approach in primary care. *CMAJ* 183, E959-
13 67.
14
15 Knipscheer, J. W., Sleijpen, M., Mooren, T., Ter Heide, F. J. & van der Aa, N. (2015). Trauma
16 exposure and refugee status as predictors of mental health outcomes in treatment-
17 seeking refugees. *BJPsych Bull* 39, 178-82.
18
19
20 Kroenke, K., Spitzer, R. L. & Williams, J. B. (2001). The PHQ-9: validity of a brief depression
21 severity measure. *Journal of General Internal Medicine* 16, 606-13.
22
23 Laban, C.J., Gernaat, H.B., Komproe, I.H., Schreuders, B.A. & De Jong, J.T. (2004). Impact of a
24 long asylum procedure on the prevalence of psychiatric disorders in Iraqi asylum seekers
25 in The Netherlands. *Journal of Nervous and Mental Disorders* 192, 843-51.
26
27
28 Mesa-Vieira, C., Haas, A. D., Buitrago-Garcia, D., Roa-Diaz, Z. M., Minder, B., Gamba, M.,
29 Salvador, D., Jr., Gomez, D., Lewis, M., Gonzalez-Jaramillo, W. C., Pahud de Mortanges, A.,
30 Buttia, C., Muka, T., Trujillo, N. & Franco, O. H. (2022). Mental health of migrants with pre-
31 migration exposure to armed conflict: a systematic review and meta-analysis. *Lancet*
32 *Public Health* 7, e469-e481.
33
34 Miller, K. E., Jordans, M. J. D., Tol, W. A. & Galappatti, A. (2021). A call for greater conceptual
35 clarity in the field of mental health and psychosocial support in humanitarian settings.
36 *Epidemiology and Psychiatric Sciences* 30, e5.
37
38
39 Mollica, R. F., Caspi-Yavin, Y., Bollini, P., Truong, T., Tor, S. & Lavelle, J. (1992). The Harvard
40 Trauma Questionnaire. Validating a cross-cultural instrument for measuring torture,
41 trauma, and posttraumatic stress disorder in Indochinese refugees. *Journal of Nervous*
42 *and Mental Disorders* 180, 111-6.
43
44
45 Nickerson, A., Byrow, Y., Rasmussen, A., O'Donnell, M., Bryant, R., Murphy, S., Mau, V.,
46 McMahon, T., Benson, G. & Liddell, B. (2021). Profiles of exposure to potentially traumatic
47 events in refugees living in Australia. *Epidemiology and Psychiatric Sciences* 30, e18.
48
49
50 Pandi-Perumal, S. R., Kumar, V. M., Pandian, N. G., de Jong, J. T., Andiappan, S., Corlățeanu, A.,
51 Mahalaksmi, A. M., Chidambaram, S. B., Kumar, R. R., Ramasubramanian, C.,
52 Sivasubramaniam, S., Bjorkum, A. A., Cutajar, J., Berk, M., Trakht, I., Vrdoljak, A., Meira, E.
53 C. M., Eyre, H. A., Gronli, J., Cardinali, D. P., Maercker, A., van de Put, W., Guzder, J.,
54 Bjorvatn, B., Tol, W. A., Acuna-Castroviejo, D., Meudec, M., Morin, C. M., Partinen, M.,
55 Barbui, C., Jordans, M. J. D., Braakman, M. H., Knaevelsrud, C., Pallesen, S., Sijbrandij, M.,
56 Golombek, D. A., Espie, C. A., Cuijpers, P., Agudelo, H. A. M., van der Velden, K., van der
57 Kolk, B. A., Hobfoll, S. E., Deville, W., Gradisar, M., Riemann, D., Axelsson, J., Benitez-King,
58 G., Macy, R. D., Poberezhets, V., Hoole, S. R. H., Murthy, R. S., Hegemann, T., Heinz, A.,
59 Salvage, J., McFarlane, A. C., Keukens, R., de Silva, H., Oestereich, C., Wilhelm, J., von
60

- 1
2
3
4 Cranach, M., Hoffmann, K., Klosinski, M., Bhugra, D. & Seeman, M. V. (2022). Scientists
5 Against War: A Plea to World Leaders for Better Governance. *Sleep and Vigilance*, 1-6.
6
7 Papola, D., Purgato, M., Gastaldon, C., Bovo, C., van Ommeren, M., Barbui, C. & Tol, W. A. (2020).
8 Psychological and social interventions for the prevention of mental disorders in people
9 living in low- and middle-income countries affected by humanitarian crises. *Cochrane*
10 *Database of Systematic Reviews* 9, CD012417.
11
12 Patel, A. R., Newman, E. & Richardson, J. (2022). A pilot study adapting and validating the
13 Harvard Trauma Questionnaire (HTQ) and PTSD checklist-5 (PCL-5) with Indian women
14 from slums reporting gender-based violence. *BMC Womens Health* 22, 22.
15
16 Plumpton, C. O., Morris, T., Hughes, D. A. & White, I. R. (2016). Multiple imputation of multiple
17 multi-item scales when a full imputation model is infeasible. *BMC Research Notes* 9, 45.
18
19 Priebe, S., Giacco, D. & El-Nagib, R. (2016). *Public Health Aspects of Mental Health Among*
20 *Migrants and Refugees: A Review of the Evidence on Mental Health Care for Refugees,*
21 *Asylum Seekers and Irregular Migrants in the WHO European Region.* WHO Regional
22 Office for Europe: Copenhagen.
23
24 Purgato, M., Carswell, K., Acarturk, C., Au, T., Akbai, S., Anttila, M., Baumgartner, J., Bailey, D.,
25 Biondi, M., Bird, M., Churchill, R., Eskici, S., Hansen, L. J., Heron, P., Ilkkursun, Z., Kilian, R.,
26 Koesters, M., Lantta, T., Nose, M., Ostuzzi, G., Papola, D., Popa, M., Sijbrandij, M.,
27 Tarsitani, L., Tedeschi, F., Turrini, G., Uygun, E., Valimaki, M. A., Wancata, J., White, R.,
28 Zanini, E., Cuijpers, P., Barbui, C. & Van Ommeren, M. (2019). Effectiveness and cost-
29 effectiveness of Self-Help Plus (SH+) for preventing mental disorders in refugees and
30 asylum seekers in Europe and Turkey: study protocols for two randomised controlled
31 trials. *BMJ Open* 9, e030259.
32
33 Purgato, M., Carswell, K., Tedeschi, F., Acarturk, C., Anttila, M., Au, T., Bajbouj, M., Baumgartner,
34 J., Biondi, M., Churchill, R., Cuijpers, P., Koesters, M., Gastaldon, C., Ilkkursun, Z., Lantta,
35 T., Nose, M., Ostuzzi, G., Papola, D., Popa, M., Roselli, V., Sijbrandij, M., Tarsitani, L.,
36 Turrini, G., Valimaki, M., Walker, L., Wancata, J., Zanini, E., White, R., van Ommeren, M. &
37 Barbui, C. (2021). Effectiveness of Self-Help Plus in Preventing Mental Disorders in
38 Refugees and Asylum Seekers in Western Europe: A Multinational Randomized Controlled
39 Trial. *Psychotherapy and Psychosomatics* 90, 403-414.
40
41 Royston, P. (2005). Multiple imputation of missing values: update. *Stata Journal* 5.
42
43 Rytwinski, N. K., Scur, M. D., Feeny, N. C. & Youngstrom, E. A. (2013). The co-occurrence of major
44 depressive disorder among individuals with posttraumatic stress disorder: a meta-
45 analysis. *Journal of Traumatic Stress* 26, 299-309.
46
47 Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T.,
48 Baker, R. & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview
49 (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview
50 for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* 59 Suppl 20, 22-33;quiz 34-57.
51
52 Statacorp (2017). Statacorp.
53
54
55
56
57
58
59
60

- 1
2
3
4 Sterne, J. A., White, I. R., Carlin, J. B., Spratt, M., Royston, P., Kenward, M. G., Wood, A. M. &
5 Carpenter, J. R. (2009). Multiple imputation for missing data in epidemiological and clinical
6 research: potential and pitfalls. *BMJ* 338, b2393.
7
8
9 Stevens, J. (2002). *Applied multivariate statistics for the social sciences*. Lawrence Erlbaum
10 Associates: Mahwah, N.J.
11
12 Tabachnick, B. F., LS (2007). *Using multivariate statistics*. Pearson/Allyn & Bacon: Boston.
13
14 Turrini, G., Tedeschi, F., Cuijpers, P., Del Giovane, C., Kip, A., Morina, N., Nose, M., Ostuzzi, G.,
15 Purgato, M., Ricciardi, C., Sijbrandij, M., Tol, W. & Barbui, C. (2021). A network meta-
16 analysis of psychosocial interventions for refugees and asylum seekers with PTSD. *BMJ*
17 *Global Health* 6.
18
19 UNHCR (2022). Refugee data finder. United Nations High Commissioner for Refugees.
20
21 Uphoff, E., Robertson, L., Cabieses, B., Villalon, F. J., Purgato, M., Churchill, R. & Barbui, C. (2020).
22 An overview of systematic reviews on mental health promotion, prevention, and
23 treatment of common mental disorders for refugees, asylum seekers, and internally
24 displaced persons. *Cochrane Database of Systematic Reviews* 9, CD013458.
25
26
27 White, I. R., Royston, P. & Wood, A. M. (2011). Multiple imputation using chained equations:
28 Issues and guidance for practice. *Statistics in Medicine* 30, 377-99.
29
30
31 White, R. G. & Van der Boor, C. (2021). Enhancing the capabilities of forcibly displaced people: a
32 human development approach to conflict- and displacement-related stressors.
33 *Epidemiology and Psychiatric Sciences* 30, e34.
34
35 World Health Organization, W. H. (2017). *Scalable psychological interventions for people in*
36 *communities affected by adversity*. World Health Organization: Geneva.
37
38 World Health Organization, W. H. (2021a). Self Help Plus (SH+): a group-based stress
39 management course for adults. Generic field-trial version 1.0. World Health Organization:
40 Geneva.
41
42 World Health Organization, W. H. (2021b). *Doing What Matters in Times of Stress*. World Health
43 Organization: Geneva.
44
45 Winkler, J.G., Brandl, E.J., Bretz, H.J., Heinz, A. & Schouler-Ocak, M. (2019) The Influence of
46 Residence Status on Psychiatric Symptom Load of Asylum Seekers in Germany.
47 *Psychiatrische Praxis* 46, 191-199
48
49
50 Wu, Y., Levis, B., Sun, Y., Krishnan, A., He, C., Riehm, K. E., Rice, D. B., Azar, M., Yan, X. W.,
51 Neupane, D., Bhandari, P. M., Imran, M., Chiovitti, M. J., Saadat, N., Boruff, J. T., Cuijpers,
52 P., Gilbody, S., McMillan, D., Ioannidis, J. P. A., Kloda, L. A., Patten, S. B., Shrier, I.,
53 Ziegelstein, R. C., Henry, M., Ismail, Z., Loiselle, C. G., Mitchell, N. D., Tonelli, M., Al-Adawi,
54 S., Beraldi, A., Braeken, A., Buel-Drabe, N., Bunevicius, A., Carter, G., Chen, C. K., Cheung,
55 G., Clover, K., Conroy, R. M., Cukor, D., da Rocha, E. S. C. E., Dabscheck, E., Daray, F. M.,
56 Douven, E., Downing, M. G., Feinstein, A., Ferentinos, P. P., Fischer, F. H., Flint, A. J.,
57 Fujimori, M., Gallagher, P., Gandy, M., Goebel, S., Grassi, L., Harter, M., Jenewein, J., Jette,
58 N., Juliao, M., Kim, J. M., Kim, S. W., Kjaergaard, M., Kohler, S., Loosman, W. L., Lowe, B.,
59
60

1
2
3
4 Martin-Santos, R., Massardo, L., Matsuoka, Y., Mehnert, A., Michopoulos, I., Misery, L.,
5 Navines, R., O'Donnell, M. L., Ozturk, A., Peceliuniene, J., Pintor, L., Ponsford, J. L., Quinn,
6 T. J., Reme, S. E., Reuter, K., Rooney, A. G., Sanchez-Gonzalez, R., Schwarzbald, M. L.,
7 Senturk Cankorur, V., Shaaban, J., Sharpe, L., Sharpe, M., Simard, S., Singer, S., Stafford,
8 L., Stone, J., Sultan, S., Teixeira, A. L., Tiringer, I., Turner, A., Walker, J., Walterfang, M.,
9 Wang, L. J., White, J., Wong, D. K., Benedetti, A. & Thombs, B. D. (2020). Probability of
10 major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic
11 interviews controlling for Hospital Anxiety and Depression Scale - Depression subscale
12 scores: An individual participant data meta-analysis of 73 primary studies. *Journal of*
13 *Psychosomatics Research* 129, 109892.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1. Participants characteristics at baseline

Means	M (SD)
Age (years)	31.797 (9.521)
Years of education	9.556 (4.424)
Months in host country	40.636 (32.599)
Host country GDP in thousand dollars (2019)	
Austria	58,091.3
Finland	50,321.6
Germany	55 652.9
Italy	44,334.2
Turkey	27,144.2
UK	49,070.3
Number of SH+ sessions	1.434 (2.015)
GHQ	5.737 (2.230)
PHQ-9	7.232 (5.230)
HTQ Total score	5.795 (4.349)
HTQ Lack of basic needs	1.277 (1.210)
HTQ Violence and abuse	1.753 (2.085)
HTQ Being close to death	2.766 (1.947)
Frequencies	n/N (%)
Gender (Female)	538/1,101 (48.86%)
Unemployed	418/1,099 (38.03%)
Country of origin	
Nigeria	114/1,100 (10.36%)
Syria	758/1,100 (68.91%)
Iraq	94/1,100 (8.55%)
Other	134/1,100 (12.18%)
Travel duration	
Less than one month	277 (40.44%)
One to three months	203 (29.64%)
More than three months	205 (29.93%)
Study	
Western Europe	459 (41.69%)
Turkey	642 (58.31%)

M: mean; SD: standard deviation; SH+: Self-Help Plus; GDP: Gross Domestic Product; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ: Harvard Trauma Questionnaire

Table 2. Determinants of mental disorder development: logistic regression analyses

	Unadjusted regressions		Adjusted regressions			
			With HTQ total score		With HTQ factors	
	Odds Ratio (95% CI)	P-value	Odds Ratio (95% CI)	P-value	Odds Ratio (95% CI)	P-value
Age in years	1.001 (0.986; 1.015)	0.920	0.996 (0.980; 1.013)	0.672	0.998 (0.981; 1.015)	0.830
Female gender	1.318 (0.987; 1.759)	0.061	0.962 (0.680; 1.360)	0.825	0.977 (0.689; 1.385)	0.896
Years of education	0.972 (0.941; 1.004)	0.088	0.959 (0.923; 0.997)	0.034	0.959 (0.922; 0.996)	0.032
Unemployed	0.938 (0.702; 1.253)	0.664	1.171 (0.832; 1.647)	0.365	1.180 (0.836; 1.664)	0.347
Country of origin:						
Syria	Reference category		Reference category		Reference category	
Iraq	1.180 (0.710; 1.960)	<0.001	1.592 (0.807; 3.142)	0.017	1.672 (0.840; 3.325)	0.006
Nigeria	0.209 (0.101; 0.432)		0.376 (0.142; 0.997)		0.318 (0.118; 0.861)	
Other	0.403 (0.240; 0.674)		0.577 (0.273; 1.222)		0.519 (0.241; 1.114)	
Travel duration (months)	Reference category				Reference category	
Less than one month	0.616 (0.398; 0.952)	<0.001	0.653 (0.414; 1.028)	0.057	0.658 (0.417; 1.041)	0.057
One to three months	0.312 (0.198; 0.494)		0.512 (0.267; 0.983)		0.502 (0.260; 0.969)	
More than three months						
Months in host country	1.002 (0.997; 1.007)	0.503	1.000 (0.994; 1.006)	0.943	1.000 (0.994; 1.006)	0.973
Turkish trial	2.058 (1.513; 2.800)	<0.001	1.795 (0.997; 3.235)	0.051	1.802 (0.998; 3.253)	0.051
Number of SH+ sessions	0.888 (0.827; 0.953)	0.001	0.853 (0.788; 0.923)	<0.001	0.850 (0.785; 0.921)	<0.001

GHQ-12	1.188 (1.114; 1.268)	<0.001	1.117 (1.036; 1.204)	0.004	1.119 (1.038; 1.206)	0.003
PHQ-9	1.071 (1.042; 1.100)	<0.001	1.061 (1.026; 1.097)	0.001	1.061 (1.026; 1.097)	0.001
HTQ Total score	1.002 (0.970; 1.036)	0.883	1.051 (1.008; 1.096)	0.020	-	-
HTQ Factor 1 - Lack of basic needs	0.999 (0.890; 1.122)	0.988	-	-	1.007 (0.863; 1.176)	0.928
HTQ Factor 2 - Violence and abuse	0.995 (0.929; 1.066)	0.894	-	-	1.163 (1.041; 1.299)	0.008
HTQ Factor 3 - Being close to death	1.018 (0.947; 1.095)	0.769	-	-	0.978 (0.888; 1.077)	0.650

CI: Confidence Interval; SH+: Self-Help Plus; GDP: Gross Domestic Product; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ: Harvard Trauma Questionnaire

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46




Figure 1. Flow-diagram of participant enrolment and follow-up




Figure 2. Frequency of mental disorders, as measured with the M.I.N.I., after three and six months of follow-up

1
2
3
4 **Risk factors for mental disorder development in asylum seekers and**
5 **refugees resettled in Western Europe and Turkey: participant-level**
6 **analysis of two large prevention trials**
7
8
9

10
11
12 **Supplemental Materials**
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Factor loadings from principal component analysis of HTQ items

	Factor 1	Factor 2	Factor 3
Lack of food or water	-0.031	-0.031	0.952
Ill health w/o access to medical care	0.057	0.037	0.841
Lack of shelter	0.155	0.079	0.745
Imprisonment	0.593	0.205	0.144
Serious injury	0.475	0.249	0.267
Combat situation	-0.374	0.590	0.315
Brain washing	0.673	-0.151	0.206
Rape or sexual abuse	0.871	0.018	-0.114
Forced isolation from others	0.706	0.085	0.180
Being close to death	0.297	0.460	0.276
Forced separation from family	0.547	0.304	0.095
Murder of family or friends	0.066	0.779	0.058
Unnatural death of family or friends	0.053	0.806	0.082
Murder of stranger(s)	-0.028	0.906	-0.039
Lost or kidnapped	0.372	0.657	-0.061
Torture	0.671	0.366	0.040
Other (e.g. domestic violence)	0.659	-0.260	0.266

HTQ: Harvard Trauma Questionnaire

Factor loadings above 0.4 marked in bold

1
2
3
4 **Secondary analysis: logistic regression including only the enhanced care as usual (ECAU) arm**
5
6
7

8
9

	Odds Ratio	95% CI	P-value
Age in years	0.998	0.975; 1.021	0.849
Female gender	0.935	0.595; 1.469	0.755
Unemployed	1.101	0.694; 1.747	0.611
Years of education	0.962	0.915; 1.011	0.126
Months in host country	0.996	0.985; 1.008	0.608
Turkish trial	1.912	0.876; 4.173	0.081
Country of origin:			
Syria	Reference cat.		
Iraq	1.225	0.501; 2.998	0.415
Nigeria	0.412	0.118; 1.440	
Other	0.691	0.263; 1.821	
Travel duration (months)			
Less than one month	Reference cat.		
One to three months	0.611	0.323; 1.156	0.071
More than three months	0.412	0.161; 1.054	
GHQ-12	1.131	1.024; 1.249	0.018
PHQ-9	1.048	1.001; 1.097	0.045
HTQ Total score	1.067	1.002; 1.136	0.037

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40
41 CI: Confidence Interval; GDP: GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ:
42 Harvard Trauma Questionnaire

43
44 Significant values marked in bold
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Secondary analysis: logistic regression including only the study conducted in Western Europe**
5
6
7

	Odds Ratio	95% CI	P-value
Age in years	0.979	0.949; 1.009	0.172
Female gender	1.093	0.555; 2.154	0.796
Unemployed	1.448	0.748; 2.804	0.271
Years of education	0.986	0.927; 1.048	0.646
Number of SH+ sessions	0.903	0.776; 1.052	0.190
Months in host country	0.996	0.979; 1.014	0.669
Country of origin:			
Syria	Reference cat.		
Iraq	1.571	0.675; 3.652	0.004
Nigeria	0.204	0.065; 0.640	
Other	0.388	0.151; 0.999	
Travel duration (months)			
Less than one month	Reference cat.		
One to three months	0.990	0.431; 2.276	0.589
More than three months	0.668	0.245; 1.822	
GHQ-12	1.114	0.972; 1.276	0.120
PHQ-9	1.088	1.026; 1.154	0.005
HTQ Total score	1.091	1.007; 1.183	0.034

CI: Confidence Interval; SH+: Self-Help Plus; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ: Harvard Trauma Questionnaire

Significant values marked in bold

Secondary analysis: logistic regression including only the study conducted in Turkey

	Odds Ratio	95% CI	P-value
Age in years	1.005	0.984; 1.026	0.647
Female gender	0.942	0.618; 1.437	0.782
Unemployed	1.087	0.713; 1.657	0.698
Years of education	0.935	0.887; 0.986	0.014
Number of SH+ sessions	0.821	0.745; 0.905	<0.001
Months in host country	1.003	0.996; 1.010	0.474
Country of origin: Syria	0.225	0.061; 0.826	0.025
Travel duration (months)			
Less than one month	Reference cat.		
One to three months	0.577	0.308; 1.1080	0.160
More than three months	0.661	0.112; 3.899	
GHQ-12	1.092	0.995; 1.198	0.065
PHQ-9	1.052	1.006; 1.101	0.028
HTQ Total score	1.056	1.001; 1.114	0.045

CI: Confidence Interval; SH+: Self-Help Plus; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; HTQ: Harvard Trauma Questionnaire

1
2
3
4 **Secondary analysis: logistic regression including recruiting centres as independent variable**
5
6
7

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

	Odds Ratio	95% CI	P-value
Age in years	0.997	0.980; 1.014	0.739
Female gender	0.967	0.683; 1.369	0.850
Unemployed	1.117	0.790; 1.578	0.532
Years of education	0.951	0.914; 0.990	0.015
Number of SH+ sessions	0.851	0.786; 0.922	<0.001
Months in host country	1.001	0.995; 1.007	0.856
Country of origin:			
Syria	Reference cat.		
Iraq	1.782	0.800; 3.969	0.090
Nigeria	0.388	0.080; 1.872	
Other	1.181	0.354; 3.934	
Travel duration (months)			
Less than one month	Reference cat.		
One to three months	0.672	0.420; 1.076	0.039
More than three months	0.445	0.226; 0.874	
Recruitment centre			
Istanbul	Reference cat.		
Wien	0.093	0.017; 0.512	
Liverpool	0.399	0.156; 1.022	0.030
Turku	0.469	0.200; 1.100	
Ulm	0.880	0.355; 2.179	
Verona	0.602	0.137; 2.646	
York	0.720	0.270; 1.915	
GHQ-12	1.108	1.026; 1.195	0.009
PHQ-9	1.064	1.029; 1.101	<0.001
HTQ Total score	1.047	1.003; 1.093	0.035

51 CI: Confidence Interval; SH+: Self-Help Plus; GHQ: General Health Questionnaire; PHQ: Patient Health
52 Questionnaire; HTQ: Harvard Trauma Questionnaire

53
54 Significant values marked in bold
55
56
57
58
59
60